



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

September 09, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 10/651,604
FILING DATE: *August 29, 2003*

Certified by



Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office



BEST AVAILABLE COPY

15992 U.S. PTO
08/29/03

Certificate of Mailing

Date of Deposit: August 29, 2003 Label Number: EV247566292US
 I hereby certify under 37 C.F.R. § 1.10 that this correspondence is being deposited with the United States Postal Service as
 "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to Mail Stop Patent
 Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.


Guy Beardsley
 Printed name of person mailing correspondence

Guy Beardsley
 Signature of person mailing correspondence

19704 U.S. PTO
10/651604
08/29/03

UTILITY PATENT APPLICATION TRANSMITTAL UNDER 37 C.F.R. § 1.53(b)

Attorney Docket Number	50304/012001
Applicant	Mark Jozef Albert WAER et al.
Title	IMMUNOSUPPRESSIVE EFFECTS OF PTERIDINE DERIVATIVES
PRIORITY INFORMATION:	
This application is a continuation-in-part of pending U.S. Application No. 09/869,468, filed October 10, 2001, which is the National Stage of International Application No. PCT/EP99/10320, filed December 28, 1999, which was published in English under PCT Article 21(2), and which claims the benefit of U.S. provisional application No. 60/113,989 filed December 28, 1998; the disclosures of which are incorporated by reference in their entirety.	
SMALL ENTITY STATUS:	
<input checked="" type="checkbox"/> Applicant claims small entity status under 37 C.F.R. § 1.27.	
APPLICATION ELEMENTS:	
Cover sheet	1 page
Specification	86 pages
Claims	26 pages
Abstract	1 page
Drawings	5 sheets
Combined Declaration and Power of Attorney, which is: <input checked="" type="checkbox"/> Unsigned; <input type="checkbox"/> Newly signed for this application; <input type="checkbox"/> A copy from prior application [**SERIAL NUMBER**] and the entire disclosure of the prior application is considered as being part of the disclosure of this new application and is hereby incorporated by reference therein.	2 pages
Sequence Statement	[**] pages
Sequence Listing on Paper	[**] pages
Sequence Listing on Diskette	[**] disk
Preliminary Amendment	[**] pages
Information Disclosure Statement	[**] pages
Form PTO 1449	[**] pages

Cited References	[**] references
Recordation Form Cover Sheet and Assignment	[**] pages
English Translation	[**] pages
Certified Copy of Priority Document	[**] pages
Non-publication Request under 35 U.S.C. § 122(b).	[**] pages
Request for Deferral of Examination under 37 C.F.R. § 1.103(d)	[**] pages
A Small Entity Statement	[**] pages
Return Receipt Postcard	1
FILING FEES:	
Basic Filing Fee: \$375	\$375.00
Excess Claims Fee: 22 - 20 x 9	\$18.00
Excess Independent Claims Fee: 3 - 3 x \$42	\$0
Multiple Dependent Claims Fee: \$280/\$140	
Total Fees:	\$393.00
<input checked="" type="checkbox"/> Enclosed is a check for \$393.00 to cover the total fees. <input type="checkbox"/> Charge [**AMOUNT**] to Deposit Account No. 03-2095 to cover the total fees. <input type="checkbox"/> The filing fee is not being paid at this time. <input checked="" type="checkbox"/> Please apply any other charges or any credits to Deposit Account No. 03-2095.	
CORRESPONDENCE ADDRESS:	
James D. DeCamp, Ph.D. Reg. No. 43,580 Clark & Elbing LLP 101 Federal Street Boston, MA 02110 CUSTOMER NO.: 21559	
<div style="display: flex; justify-content: space-between;"> <div>  James D. DeCamp, Ph.D. Reg. No. 43,580 </div> <div> Telephone: 617-428-0200 Facsimile: 617-428-7045 </div> </div>	
<div style="display: flex; justify-content: space-between;"> <div></div> <div> <u>29 August 2003</u> Date </div> </div>	

Certificate of Mailing

Date of Deposit: August 29, 2003

Label Number: EV247566292US

I hereby certify under 37 C.F.R. § 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to Mail Stop Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Guy Beardsley
Printed name of person mailing correspondence


Signature of person mailing correspondence

APPLICATION
FOR
UNITED STATES LETTERS PATENT

APPLICANT : MARK JOZEF ALBERT WAER, PIET ANDRÉ
MAURITS MARIA HERDEWIJN, AND
WOLFGANG EUGEN PFLEIDERER

TITLE : IMMUNOSUPPRESSIVE EFFECTS OF
PTERIDINE DERIVATIVES

IMMUNOSUPPRESSIVE EFFECTS OF PTERIDINE DERIVATIVES

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of pending U.S. Application No. 09/869,468, filed October 10, 2001, which is the National Stage of
5 International Application No. PCT/EP99/10320, filed December 28, 1999, which was published in English under PCT Article 21(2), and which claims the benefit of U.S. provisional application No. 60/113,989 filed December 28, 1998; the disclosures of which are incorporated by reference in their entirety.

10 The invention relates to a class of novel pteridines. The invention further relates to pharmaceutical compositions including a broad class of pteridines especially for the prevention and/or the treatment of pathologic conditions such as, but not limited to, immune and auto-immune disorders, organ and cells transplant rejections, cell proliferative disorders, cardiovascular
15 disorders, disorders of the central nervous system and viral diseases.

The invention further relates to combined pharmaceutical preparations comprising one or more pteridines and one or more known immunosuppressant drugs or antineoplastic drugs or anti-viral drugs.

This invention also relates to a method for the prevention and/or
20 treatment of pathologic conditions such as, but not limited to, immune and autoimmune disorders, organ and cells transplant rejections, cell proliferative disorders, cardiovascular disorders, disorders of the central nervous system and viral diseases by the administration of an effective amount of a specific pteridine optionally combined with one or more known immunosuppressant
25 drugs or antineoplastic drugs or anti-viral drugs.

BACKGROUND OF THE INVENTION

Several 2,4-diaminopteridine derivatives being substituted in the 6-position and/or the 7-position of the pteridine ring (according to standard atom
30 numbering for said ring) are known in the art, e.g. from various sources of literature including Swiss Patent No. 231,852; British Patent No. 763,044; United States Patents No. 2,512,572; No. 2,581,889; No. 2,665,275; No. 2,667,486; No.

2,940,972; No. 3,081,230 and No. 5,047,405. Some of these substituted 2,4-diaminopteridine derivatives were disclosed in relationship with various medical uses, such as bacterial growth inhibitors, antineoplastic agents, anti-schistosomiasis activity, coronary dilating activity, diuretic and hypotensive activity, and anti-amnesic activity. In particular, U.S. Patent No. 2,940,972 and EP-A-362,645 disclose very specific 2,4-diaminopteridine derivatives being substituted by piperidinyl, morpholinyl or pyrrolidinyl in the 7-position of the pteridine ring.

Specific 2-aminopteridine derivatives wherein the 4-position of the pteridine ring is substituted with an alkoxy group and the 6-position is also substituted are also known in the art, although without any medical utility. For instance, United States Patent No. 2,740,784 discloses such derivatives wherein the 6-substituent is an acetal; J. Chem. Soc. (1957) 2146-2158 discloses 2-dimethylamino-4-ethoxy-6-phenylpteridine as a compound with a melting point of 200°C; Arm. Khim. J. discloses 2-amino-4-ethoxy-6,7-diphenylpteridine; Helv. Chem. Acta (1992) 75:2317-2326 discloses 2-amino-4-pentoxy-6-methylthiopteridine. Furthermore, International Patent Application published as WO 01/19825 discloses 2-phenylamino and 2-phenylsulfoxide pteridines wherein the 7-position is substituted with an amide as being useful inhibitors of cell cycle regulatory kinases.

Nevertheless, there still is a need in the art for specific and highly therapeutically active compounds, such as, but not limited to, drugs for treating immune and autoimmune disorders, organ and cells transplant rejections, cell proliferative disorders, cardiovascular disorders, disorders of the central nervous system and viral diseases. In particular, there is a need in the art to provide immunosuppressive compounds or antineoplastic drugs or anti-viral drugs which are active in a minor dose in order to replace existing drugs having significant side effects and to decrease treatment costs.

Currently used immunosuppressive drugs include antiproliferative agents, such as methotrexate (a 2,4-diaminopteridine derivative disclosed by U.S. Patent No. 2,512,572), azathioprine, and cyclophosphamide. Since these drugs affect mitosis and cell division, they have severe toxic effects on normal cells with high turn-over rate such as bone marrow cells and the

gastrointestinal tract lining. Accordingly, marrow depression and liver damage are common side effects of these antiproliferative drugs.

Anti-inflammatory compounds used to induce immunosuppression include adrenocortical steroids such as dexamethasone and prednisolone.

- 5 The common side effects observed with the use of these compounds are frequent infections, abnormal metabolism, hypertension, and diabetes.

- Other immunosuppressive compounds currently used to inhibit lymphocyte activation and subsequent proliferation include cyclosporine, tacrolimus and rapamycin. Cyclosporine and its relatives are among the most
10 commonly used immunosuppressant drugs. Cyclosporine is typically used for preventing or treating organ rejection in kidney, liver, heart, pancreas, bone marrow, and heart-lung transplants, as well as for the treatment of autoimmune and inflammatory diseases such as Crohn's disease, aplastic anemia, multiple-sclerosis, myasthenia gravis, uveitis, biliary cirrhosis, etc.
15 However, cyclosporines suffer from a small therapeutic dose window and severe toxic effects including nephrotoxicity, hepatotoxicity, hypertension, hirsutism, cancer, and neurotoxicity.

- Additionally, monoclonal antibodies with immunosuppressant properties, such as OKT3, have been used to prevent and/or treat graft
20 rejection. Introduction of such monoclonal antibodies into a patient, as with many biological materials, induces several side-effects, such as dyspnea. Within the context of many life-threatening diseases, organ transplantation is considered a standard treatment and, in many cases, the only alternative to death. The immune response to foreign cell surface antigens on the graft,
25 encoded by the major histo-compatibility complex (hereinafter referred as MHC) and present on all cells, generally precludes successful transplantation of tissues and organs unless the transplant tissues come from a compatible donor and the normal immune response is suppressed. Other than identical twins, the best compatibility and thus, long term rates of engraftment, are
30 achieved using MHC identical sibling donors or MHC identical unrelated cadaver donors. However, such ideal matches are difficult to achieve. Further, with the increasing need of donor organs an increasing shortage of transplanted organs currently exists. Accordingly, xenotransplantation has

emerged as an area of intensive study, but faces many hurdles with regard to rejection within the recipient organism.

The host response to an organ allograft involves a complex series of cellular interactions among T and B lymphocytes as well as macrophages or dendritic cells that recognize and are activated by foreign antigen. Co-stimulatory factors, primarily cytokines, and specific cell-cell interactions, provided by activated accessory cells such as macrophages or dendritic cells are essential for T-cell proliferation. These macrophages and dendritic cells either directly adhere to T-cells through specific adhesion proteins or secrete cytokines that stimulate T-cells, such as IL-12 and IL-15. Accessory cell-derived co-stimulatory signals stimulate activation of interleukin-2 (IL-2) gene transcription and expression of high affinity IL-2 receptors in T-cells. IL-2 is secreted by T lymphocytes upon antigen stimulation and is required for normal immune responsiveness. IL-2 stimulates lymphoid cells to proliferate and differentiate by binding to IL-2 specific cell surface receptors (IL-2R). IL-2 also initiates helper T-cell activation of cytotoxic T-cells and stimulates secretion of interferon- γ which in turn activates cytodestructive properties of macrophages. Furthermore, IFN- γ and IL-4 are also important activators of MHC class II expression in the transplanted organ, thereby further expanding the rejection cascade by enhancing the immunogenicity of the grafted organ. The current model of a T-cell mediated response suggests that T-cells are primed in the T-cell zone of secondary lymphoid organs, primarily by dendritic cells. The initial interaction requires cell to cell contact between antigen-loaded MHC molecules on antigen-presenting cells (hereinafter referred as APC) and the T-cell receptor/CD3 complex on T-cells. Engagement of the TCR/CD3 complex induces CD154 expression predominantly on CD4 T-cells that in turn activate the APC through CD40 engagement, leading to improved antigen presentation. This is caused partly by upregulation of CD80 and CD86 expression on the APC, both of which are ligands for the important CD28 co-stimulatory molecule on T-cells. However, engagement of CD40 also leads to prolonged surface expression of MHC-antigen complexes, expression of ligands for 4-1BB and OX-40 (potent co-stimulatory molecules expressed on activated T-cells). Furthermore, CD40 engagement leads to secretion of

various cytokines (e.g., IL-12, IL-15, TNF- α , IL-1, IL-6, and IL-8) and chemokines, all of which have important effects on both APC and T-cell activation and maturation. Similar mechanisms are involved in the development of auto-immune disease, such as type I diabetes. In humans and
5 non-obese diabetic mice, insulin-dependent diabetes mellitus results from a spontaneous T-cell dependent auto-immune destruction of insulin-producing pancreatic β cells that intensifies with age. The process is preceded by infiltration of the islets with mononuclear cells (insulitis), primarily composed of T lymphocytes. A delicate balance between auto-aggressive T-cells and
10 suppressor-type immune phenomena determines whether expression of auto-immunity is limited to insulitis or not. Therapeutic strategies that target T-cells have been successful in preventing further progress of the auto-immune disease. These include neonatal thymectomy, administration of cyclosporine, and infusion of anti-pan T-cell, anti-CD4, or anti-CD25 (IL-2R) monoclonal
15 antibodies. The aim of all rejection prevention and auto-immunity reversal strategies is to suppress the patient's immune reactivity to the antigenic tissue or agent, with a minimum of morbidity and mortality. Accordingly, a number of drugs are currently being used or investigated for their immunosuppressive properties. As discussed above, the most commonly used
20 immunosuppressant is cyclosporine, which however has numerous side effects. Accordingly, in view of the relatively few choices for agents effective at immunosuppression with low toxicity profiles and manageable side effects, there exists a need in the art for identification of alternative immunosuppressive agents and for agents acting as complement to
25 calcineurin inhibition.

The metastasis of cancer cells represents the primary source of clinical morbidity and mortality in the large majority of solid tumors. Metastasis of cancer cells may result from the entry of tumor cells into either lymphatic or blood vessels. Invasion of lymphatic vessels results in metastasis to regional
30 draining lymph nodes. From the lymph nodes, melanoma cells for example tend to metastasize to the lung, liver, and brain. For several solid tumors, including melanoma, the absence or the presence of lymph nodes metastasis is the best predictor of patient survival. Presently, to our knowledge, no

treatment is capable of preventing or significantly reducing metastasis. Hence, there is a need in the art for compounds having such anti-metastasis effect for a suitable treatment of cancer patients.


In the field of allergy, IgE is well known for inducing allergy mainly by stimulating mast cells to release histamine. Also, asthma, being characterized by inflammation of airway and bronchospasm, is mainly induced by Th2 cytokines such as IL-5, IL-10 or IL-13. Therefore there is a need in the art for compounds that efficiently inhibit the release of these Th2 cytokines.

There is also a need in the art to improve therapeutic efficiency by providing pharmaceutical compositions or combined preparations exhibiting a synergistic effect as a result of combining two or more immunosuppressant drugs, or antineoplastic drugs or anti-viral drugs or anti-histamine drugs.

Meeting these various needs in the art constitutes the main goal of the present invention.

SUMMARY OF THE INVENTION

In a first embodiment, the present invention relates to a group of novel pteridine derivatives having the general formula (I):



wherein X represents an oxygen atom or a group with the formula $S(O)_m$ wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R_1 is a group selected from the group consisting of C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-7} alkenyl, C_{2-7} alkynyl, halo C_{1-4} alkyl, C_{3-10} cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C_{1-7} alkyl, thio C_{3-10} cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio,

- formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof,
- 5 carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkyl-amino, hydroxylalkylamino, mercaptoalkyl-amino, heterocyclic amino, hydrazino, alkylhydrazino and phenyl-hydrazino; or R_1 is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;
- 10 - Z is a group independently defined as R_1 or Z is hydrogen or the group NZ together with R_1 is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R_2 is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl, ureido; thio-ureido,
- 15 sulfonamido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino; alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C_{3-7} alkylamino; arylamino; arylalkyl-amino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C_{1-7} alkoxy; C_{3-10}
- 20 cycloalkoxy; thio C_{1-7} alkyl; arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C_{3-10} cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals;
- R_4 is an atom or a group selected from the group consisting of hydrogen; halogen; C_{1-7} alkyl; C_{2-7} alkenyl; C_{2-7} alkynyl; halo C_{1-7} alkyl;
- 25 carboxy C_{1-7} alkyl; carboxyaryl; C_{1-7} alkoxy; C_{3-10} cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C_{1-7} alkyl; thio C_{3-10} cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-
- 30 acetal; carboxylic acid; carboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido;

thio-ureido; alkylamino; cycloalkylamino; alkenylamino; cycloalkenyl-
 amino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino;
 mercaptoalkylamino; heterocyclic amino; heterocyclic-substituted
 alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino;
 5 phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides,
 amides and thioamides thereof; phenyl substituted with one or more
 substituents selected from the group consisting of C₁₋₇ alkyl, C₂₋₇
 alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino,
 C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-
 10 substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl,
 thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl,
 carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxyl-
 amino, alkoxyamino, mercaptoamino, thioalkylamino, acylamino,
 thioacylamino, cyano, carboxylic acid or esters or thioesters or halides
 15 or anhydrides or amides thereof, thiocarboxylic acid or esters or
 thioesters or halides or anhydrides or amides thereof, alkylamino,
 cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, aryl-
 amino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino,
 heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino;
 20 aryl groups other than phenyl, the said aryl groups being optionally
 substituted with one or more substituents selected from the group
 consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl,
 halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy,
 aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy,
 25 thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic,
 arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl,
 thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino,
 alkoxyamino, mercaptoamino, thioalkylamino, acylamino, thioacyl-
 amino, cyano, carboxylic acid or esters or thioesters or halides or
 30 anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters
 or halides or anhydrides or amides thereof, alkylamino, cyclo-
 alkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino,
 arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic

amino, hydrazino, alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic radicals other than piperidinyl, morpholinyl or pyrrolidinyl, i.e. preferably selected from the group consisting of

5 azacycloheptyl, azabenzimidazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydropyryl, tetrahydroquinoleinyl, tetrahydrothienyl and dioxide thereof, dihydrothienyl dioxide, dioxindolyl, dioxinyl, dioxenyl, dioxazinyl, thioxanyl, thioxolyl, thio-urazolyl, thiotriazolyl, thiopyranyl, thiopyryl, coumarinyl, quinoleinyl, oxyquinoleinyl, quinuclidinyl,

10 xanthinyl, dihydropyranyl, benzodihydrofuryl, benzothio-pyryl, benzothiopyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl, benzodioxanyl, benzothiadiaazolyl, benzotriazinyl, benzothiazolyl, benzoxazolyl, phenothioxinyl, phenothiazolyl, phenothieryl, phenopyryl, phenoxazolyl, pyridinyl, dihydropyridinyl, tetrahydro-

15 pyridinyl, piperidinyl, thiomorpholinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl, triazolyl, benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiaazolyl, isothiazolyl, oxazolyl, oxadiaazolyl, pyrrolyl, furyl, dihydrofuryl, furoyl, hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithieryl, thienyl, indolyl, indazolyl, benzofuryl,

20 quinolyl, quinazolinyl, quinoxalyl, carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothienyl, naphthothienyl, thianthrenyl, pyryl, pyryl, benzopyryl, isobenzofuranyl, chromenyl, phenoxathieryl, indolizyl, quinolizyl, isoquinolyl, phthalazinyl, naphthiridinyl, cinnolyl, pteridinyl, carbolinyl, acridinyl,

25 perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, imidazolyl, imidazolidinyl, benzimidazolyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl, pyrrolidinyl, piperazinyl, uridinyl, thymidinyl, cytidinyl, azirinyl, aziridinyl, diazirinyl, diaziridinyl, oxiranyl, oxaziridinyl, dioxiranyl, thiiranyl, azetyl,

30 dihydroazetyl, azetidyl, oxetyl, oxetanyl, thietyl, thietanyl, diaza-bicyclooctyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromanyl, chromanonyl, thiochromanyl, thiochromanonyl, thiochromenyl, benzofuranyl, benzisothiazolyl, benzocarbazolyl, benzochromonyl, benziso-alloxazinyl, benzocoumarinyl, thiocoumarinyl, phenometoxazinyl,

phenoparoxazinyl, phentriazinyl, thiodiazinyl, thiodiazolyl, indoxyl, thio-indoxyl, benzodiazinyl, phtalidyl, phtalimidinyl, phtalazonyl, alloxazinyl, xanthionyl, isatyl, isopyrazolyl, isopyrazolonyl, urazolyl, urazinyl, uretinyl, uretidinyl, succinyl, succinimido, benzylsultimyl and benzylsultamyl; aromatic or heterocyclic substituents substituted with an aliphatic spacer between the pteridine ring and the aromatic or heterocyclic substituent, whereby said aliphatic spacer is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may contain one or more functions, atoms or radicals selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 1 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, aminoacid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxy-alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; and


- R₃ is an atom or a group selected from the group consisting of fluoro, bromo, iodo, C₂₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₂₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino; acylamino; thio-acylamino; thio-acetal; carboxylic acid; carboxylic acid esters, thioesters, amides, halides, anhydrides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, amides, halides, anhydrides and thioamides; hydroxyl; sulfhydryl; nitro; carbamoyl; thiocarbamoyl; ureido; thio-ureido; amino; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercaptoalkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, amides and thioamides thereof; aryl optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, mercaptoamino, alkoxyamino, thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic radicals; aromatic or heterocyclic substituents substituted with an aliphatic spacer between the pteridine ring and the aromatic or heterocyclic substituent, whereby said aliphatic spacer is a

5 branched or straight, saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may contain one or more functions, atoms or radicals selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, carboxylic acid or ester or thioester or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, alkylamino, cycloalkylamino, alkenylamino, cyclo-alkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkyl-amino, mercaptoalkylamino, heterocyclic amino, hydrazino,
 10 alkylhydrazino, phenylhydrazino, sulfonyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 2 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid,
 15 cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thio carboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenyl-amino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino,
 20 alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; or R₃ together with R₄ forms a homocyclic or heterocyclic radical such as, but not limited to, indolyl, dihydroxypyrimidyl or tetramethylene;
 25

and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-N-oxide thereof and/or a solvate thereof and/or a dihydro- or tetrahydropteridine derivative thereof.

30 The above novel compounds have in common the structural features present in the general formula (I), in particular they are at least trisubstituted in positions 2, 4 and 6 of the pteridine ring. They also have a potential specific biological activity profile and consequent usefulness in medicinal chemistry.

In a second embodiment, the present invention relates to the unexpected finding that at least one desirable biological property such as, but not limited to, the ability to decrease the proliferation of lymphocytes, or to decrease T-cell activation, or to decrease B-cell or monocytes or macrophages activation, or to inhibit the release of certain cytokines, is a common feature which is not only present in the group of novel compounds defined in the general formula (I), but also in a group of pteridine derivatives which is broader than the said group of novel compounds. As a consequence, the invention relates to pharmaceutical compositions comprising as an active principle at least one pteridine derivative having the general formula (II):



wherein X represents an oxygen atom or a group with the formula $S(O)_m$ wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R_1 is a group selected from the group consisting of C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-7} alkenyl, C_{2-7} alkynyl, halo C_{1-4} alkyl, C_{3-10} cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C_{1-7} alkyl, thio C_{3-10} cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxylalkylamino, mercaptoalkyl-amino, heterocyclic amino, hydrazino, alkylhydrazino and phenyl-hydrazino; or R_1 is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;

- Z is a group independently defined as R₁ or Z is hydrogen or the group NZ together with R₁ is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R₂ is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl, ureido; thioureido, sulfon-amido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino; alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C₃₋₇ alkylamino; arylamino; arylalkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; thio C₁₋₇ alkyl; arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C₃₋₁₀ cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals,
- R₄ is an atom or a group selected from the group consisting of hydrogen; halogen; C₁₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; carboxy C₁₋₇ alkyl; acetoxyc C₁₋₇ alkyl; carboxyaryl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₁₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-acetal; carboxylic acid; carboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercaptoalkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and thioamides thereof, phenyl substituted with one or more substituents selected from the group consisting of C₁₋₇ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy,

arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇
 alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio,
 heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl,
 ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxyamino,
 5 mercaptoamino, thioalkylamino, acylamino, thioacylamino, cyano,
 carboxylic acid or esters or thioesters or halides or anhydrides or
 amides thereof, thiocarboxylic acid or esters or thioesters or halides or
 anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenyl-
 amino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino,
 10 hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino,
 alkylhydrazino and phenylhydrazino; aryl groups other than phenyl, the
 said aryl groups being optionally substituted with one or more
 substituents selected from the group consisting of halogen, C₁₋₇ alkyl,
 C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl,
 15 sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyhetero-
 cyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀
 cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-
 substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-
 ureido, sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino,
 20 thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or
 esters or thioesters or halides or anhydrides or amides thereof,
 thiocarboxylic acid or esters or thioesters or halides or anhydrides or
 amides thereof, alkylamino, cycloalkylamino, alkenylamino, cyclo-
 alkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkyl-
 25 amino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkyl-
 hydrazino and phenylhydrazino; optionally substituted heterocyclic
 radicals other than piperidinyl, morpholinyl or pyrrolidinyl, i.e. preferably
 selected from the group consisting of oxabicycloheptyl, azabenzimi-
 dazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl,
 30 tetrahydrofuryl, tetrahydropyranyl, tetrahydropyronyl, tetrahydroquino-
 leinyl, tetrahydrothienyl and dioxide thereof, dihydrothienyl dioxide,
 dioxindolyl, dioxinyl, dioxenyl, dioxazinyl, thioxanyl, thioxolyl, thio-
 urazolyl, thiotriazolyl, thiopyranyl, thiopyronyl, coumarinyl, quinoleinyl,

oxyquinoleinyl, quinuclidinyl, xanthinyl, dihydropyranyl, benzodihydrofuryl, benzothiopyronyl, benzothiopyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl, benzodioxanyl, benzothiadiazolyl, benzotriazinyl, benzothiazolyl, benzoxazolyl, phenothioxinyl, phenothiazolyl, phenothieryl, phenopyronyl, phenoxazolyl, pyridinyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl, thiomorpholinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl, triazolyl, benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, isothiazolyl, oxazolyl, oxadiazolyl, pyrrolyl, furyl, dihydrofuryl, furoyl, hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithiinyl, thienyl, indolyl, indazolyl, benzofuryl, quinolyl, quinazolyl, quinoxalyl, carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothieryl, naphthothieryl, thianthrenyl, pyranyl, pyronyl, benzopyronyl, isobenzofuranyl, chromenyl, phenoxathieryl, indolizyl, quinolizyl, isoquinolyl, phthalazinyl, naphthiridinyl, cinnolyl, pteridinyl, carbolyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, imidazolyl, imidazolidinyl, benzimidazolyl, pyrazolyl, pyrazolidinyl, pyrrolinyl, pyrrolidinyl, piperazinyl, uridinyl, thymidinyl, cytidinyl, azirinyl, aziridinyl, diazirinyl, diaziridinyl, oxiranyl, oxaziridinyl, dioxiranyl, thiiranyl, azetyl, dihydroazetyl, azetidyl, oxetyl, oxetanyl, thietyl, thietanyl, diaza-bicyclooctyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromanyl, chromanonyl, thiochromanyl, thiochromanonyl, thiochromenyl, benzofuranyl, benzisothiazolyl, benzocarbazolyl, benzochromonyl, benziso-alloxazinyl, benzocoumarinyl, thiocoumarinyl, phenometoxazinyl, phenoparoxazinyl, phentriazinyl, thiodiazinyl, thiodiazolyl, indoxyl, thioindoxyl, benzodiazinyl, phtalidyl, phtalimidinyl, phtalazonyl, alloxazinyl, xanthionyl, isatyl, isopyrazolyl, isopyrazolonyl, urazolyl, urazinyl, uretinyl, uretidinyl, succinyl, succinimido, benzylsultimyl and benzylsultamyl; aromatic or heterocyclic substituents substituted with an aliphatic spacer between the pteridine ring and the aromatic or heterocyclic substituent, whereby said aliphatic spacer is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may contain one or more functions, atoms or radicals selected

from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkyl-amino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 2 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyl-oximino, amino-acid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercapto-amino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxy-alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; and

- R₃ is an atom or a group defined as R₄ or R₃ is selected from the group consisting of morpholinyl, amino, hydrogen, methyl, thiomethyl and chloro; or R₃ together with R₄ forms a homocyclic or heterocyclic radical such as, but not limited to, indolyl, dihydroxypyrimidyl or tetramethylene;

and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-N-oxide thereof and/or a solvate and/or a dihydro- or tetrahydropteridine derivative thereof.

Compounds of formula (II) are highly active immunosuppressive agents, antineoplastic agents, anti-allergic agents or anti-viral agents which, together

with one or more pharmaceutically acceptable carriers, may be formulated into pharmaceutical compositions for the prevention or treatment of pathologic conditions such as, but not limited to, immune and autoimmune disorders, organ and cells transplant rejections, allergic conditions, cell proliferative disorders, cardiovascular disorders, disorders of the central nervous system and viral diseases.

In a further embodiment, the present invention relates to combined preparations containing at least one compound of formula (II) and one or more drugs such as immunosuppressant and/or immunomodulator drugs, antineoplastic drugs, anti-histamines, inhibitors of agents causative of allergic conditions, or antiviral agents. In a further embodiment, the present invention relates to the prevention or treatment of the above-cited pathologic conditions by administering to the patient in need thereof an effective amount of a compound of general formula (II), optionally in the form of a pharmaceutical composition or combined preparation with another suitable drug.

In a still further embodiment, the present invention relates to various processes and methods for making the novel pteridine derivatives defined in general formula (I), as well as their pharmaceutically acceptable salts, N-oxides, solvates, enantiomers and dihydro- and tetrahydroderivatives.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1 and 2 represent alternative schemes for preparing 2,4,6-trisubstituted pteridine derivatives according to this invention.

Figure 3 represents a scheme for preparing 2,4,7-trisubstituted pteridine derivatives according to this invention.

Figure 4 represents a scheme for preparing unsymmetrical 2,4,6-trisubstituted pteridines and 2,4,7-trisubstituted pteridines according to this invention.

Figure 5 represents a scheme for preparing symmetrical 2,4,6-trisubstituted pteridines and 2,4,7-trisubstituted pteridines according to this invention.

DEFINITIONS

Unless otherwise stated herein, the term " trisubstituted " means that three of the carbon atoms being in positions 2, 4 and 6 or, alternatively, in positions 2, 4 and 7 of the pteridine ring (according to standard atom numbering for the pteridine ring) are substituted with an atom or group other than hydrogen. The term " tetrasubstituted " means that all four carbon atoms being in positions 2, 4, 6 and 7 of the pteridine ring are substituted with an atom or group other than hydrogen.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms " C₁₋₇ alkyl" or " aliphatic saturated hydrocarbon radicals with 1 to 7 carbon atoms " means straight and branched chain saturated acyclic hydrocarbon monovalent radicals having from 1 to 7 carbon atoms such as, for example, methyl, ethyl, propyl, n-butyl, 1-methylethyl (isopropyl), 2-methylpropyl (isobutyl), 1,1-dimethylethyl (ter-butyl), 2-methylbutyl, n-pentyl, dimethylpropyl, n-hexyl, 2-methylpentyl, 3-methylpentyl, n-heptyl and the like; the term " C₁₋₄ alkyl " designate the corresponding radicals with only 1 to 4 carbon atoms, and so on.

As used herein with respect to a substituting radical, and unless otherwise stated, the term C₁₋₇ alkylene means the divalent hydrocarbon radical corresponding to the above defined C₁₋₇ alkyl, such as methylene, bis(methylene), tris(methylene), tetramethylene, hexamethylene and the like.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms " C₃₋₁₀ cycloalkyl" and " cycloaliphatic saturated hydrocarbon radical with 3 to 10 carbon atoms " means a monocyclic saturated hydrocarbon monovalent radical having from 3 to 10 carbon atoms, such as for instance cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and the like, or a C₇₋₁₀ polycyclic saturated hydrocarbon monovalent radical having from 7 to 10 carbon atoms such as, for instance, norbornyl, fenchyl, trimethyltricycloheptyl or adamantyl.

As used herein with respect to a substituting radical, and unless otherwise stated, the term " C₃₋₁₀ cycloalkylene" means the divalent hydrocarbon radical corresponding to the above defined C₃₋₁₀ cycloalkyl.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms " aryl " and "aromatic substituent " are interchangeable and designate any mono- or polyaromatic monovalent hydrocarbon radical having from 6 up to 30 carbon atoms such as but not
 5 limited to phenyl, naphthyl, anthracenyl, adamantyl, phenanthracenyl, fluoranthenyl, chrysenyl, pyrenyl, biphenyl, terphenyl, picenyl and the like, including spiro hydrocarbon radicals and fused benzo - C₅₋₈ cycloalkyl radicals (the latter being as defined above) such as, for instance, indanyl, 1,2,3,4-tetrahydronaphthalenyl, fluorenyl and the like.

10 As used herein with respect to a substituting radical such as the combination of R₃ and R₄, and unless otherwise stated, the term " homo-cyclic" means a mono- or polycyclic, saturated or mono-unsaturated or polyunsaturated hydrocarbon radical having from 4 up to 15 carbon atoms but including no heteroatom in the said ring.

15 As used herein with respect to a substituting radical, and unless otherwise stated, the term " heterocyclic " means a mono- or polycyclic, saturated or mono-unsaturated or polyunsaturated monovalent hydrocarbon radical having from 2 up to 15 carbon atoms and including one or more heteroatoms in a 3 to 10 membered ring (and optionally one or more
 20 heteroatoms attached to one or more carbon atoms of said ring, for instance in the form of a carbonyl or thiocarbonyl group) and/or to one or more heteroatoms of said ring, for instance in the form of a sulfone, sulfoxide, N-oxide, phosphate, phosphonate or selenium oxide, each said heteroatom being independently selected from the group consisting of nitrogen, oxygen,
 25 sulfur, selenium and phosphorus, including benzo-fused heterocyclic radicals, such as but not limited to oxabicycloheptyl, azabenzimidazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydropyranyl, tetrahydroquinoleinyl, tetrahydrothienyl and dioxide thereof, dihydrothienyl dioxide, dioxindolyl,
 30 dioxinyl, dioxenyl, dioxazinyl, thioxanyl, thioxolyl, thio-urazolyl, thiotriazolyl, thiopyranyl, thiopyranyl, coumarinyl, quinoleinyl, oxyquinoleinyl, quinuclidinyl, xanthinyl, dihydropyranyl, benzodihydrofuryl, benzothiopyranyl, benzothiopyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl, benzodioxanyl,

- benzothiadiazolyl, benzotriazinyl, benzothiazolyl, benzoxazolyl, phenothioxinyl, phenothiazolyl, phenothienyl (benzothiofuranyl), phenopyreryl, phenoxazolyl, pyridinyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, 5 tetrazinyl, triazolyl, benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, isothiazolyl, oxazolyl, oxadiazolyl, pyrrolyl, furyl, dihydrofuryl, furoyl, hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithiinyl, thienyl, indolyl, indazolyl, benzofuryl, quinolyl, quinazolinyl, quinoxalinyl, carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothienyl, naphthothienyl, 10 thianthrenyl, pyranyl, pyronyl, benzopyreryl, isobenzofuranyl, chromenyl, phenoxathiinyl, indoliziny, quinoliziny, isoquinolyl, phthalazinyl, naphthiridinyl, cinnoliny, pteridinyl, carbolinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, imidazoliny, imidazolidinyl, benzimidazolyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl, pyrrolidinyl, piperazinyl, uridinyl, 15 thymidinyl, cytidinyl, azirinyl, aziridinyl, diazirinyl, diaziridinyl, oxiranyl, oxaziridinyl, dioxiranyl, thiiranyl, azetyl, dihydroazetyl, azetidiny, oxetyl, oxetanyl, thietyl, thietanyl, diazabicyclo-octyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromanyl, chromanonyl, thiochromanyl, thiochromanonyl, thiochromenyl, benzofuranyl, benzisothiazolyl, benzocarbazolyl, 20 benzochromonyl, benziso-alloxazinyl, benzocoumarinyl, thiocoumarinyl, phenometoxazinyl, phenoparoxazinyl, phentriazinyl, thiodiazinyl, thiodiazolyl, indoxyl, thio-indoxyl, benzodiazinyl (e.g. phtalazinyl), phtalidyl, phtalimidinyl, phtalazonyl, alloxazinyl, dibenzopyreryl (i.e. xanthonyl), xanthionyl, isatyl, isopyrazolyl, isopyrazolonyl, urazolyl, urazinyl, uretanyl, uretidinyl, succinyl, 25 succinimido, benzylsultimyl, benzylsultamyl and the like, including all possible isomeric forms thereof, wherein each carbon atom of the said ring may be substituted with a substituent selected from the group consisting of halogen, nitro, C₁₋₇ alkyl (optionally containing one or more functions or radicals selected from the group consisting of carbonyl (oxo), alcohol (hydroxyl), ether (alkoxy), acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, 30 carboxylic acid ester or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, C₁₋₇ alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxylalkylamino, mercapto-alkylamino,

heterocyclic amino, hydrazino, alkylhydrazino, phenyl-hydrazino, sulfonyl, sulfonamido and halogen), C₃₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, C₃₋₁₀ cycloalkyl, aryl, arylalkyl, alkylaryl, alkylacyl, arylacyl, hydroxyl, amino, C₁₋₇ alkylamino, cycloalkylamino, alkenylamino, cyclo-alkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfhydryl, C₁₋₇ alkoxy, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxylamino, cyano, carboxylic acid or esters or thioesters or amides thereof, thiocarboxylic acid or esters or thioesters or amides thereof; depending upon the number of unsaturations in the 3 to 10 membered ring, heterocyclic radicals may be sub-divided into heteroaromatic (or "heteroaryl") radicals and non-aromatic heterocyclic radicals; when a heteroatom of the said non-aromatic heterocyclic radical is nitrogen, the latter may be substituted with a substituent selected from the group consisting of C₁₋₇ alkyl, C₃₋₁₀ cycloalkyl, aryl, arylalkyl and alkylaryl.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms "C₁₋₇ alkoxy", "C₃₋₁₀ cycloalkoxy", "aryloxy", "arylalkyloxy", "oxyheterocyclic", "thio C₁₋₇ alkyl", "thio C₃₋₁₀ cycloalkyl", "arylthio", "arylalkylthio" and "thioheterocyclic" refer to substituents wherein a C₁₋₇ alkyl radical, respectively a C₃₋₁₀ cycloalkyl, aryl, arylalkyl or heterocyclic radical (each of them such as defined herein), are attached to an oxygen atom or a sulfur atom through a single bond, such as but not limited to methoxy, ethoxy, propoxy, butoxy, thioethyl, thiomethyl, phenyloxy, benzyloxy, mercaptobenzyl, cresoxy and the like.

As used herein with respect to a substituting atom, and unless otherwise stated, the term halogen means any atom selected from the group consisting of fluorine, chlorine, bromine and iodine.

As used herein with respect to a substituting radical, and unless otherwise stated, the term "halo C₁₋₇ alkyl" means a C₁₋₇ alkyl radical (such as above defined) in which one or more hydrogen atoms are independently replaced by one or more halogens (preferably fluorine, chlorine or bromine),

such as but not limited to difluoromethyl, trifluoromethyl, trifluoroethyl, octafluoropentyl, dodecafluoroheptyl, dichloromethyl and the like; the term "halo C₁₋₄ alkyl " designate the corresponding radical with only 1 to 4 carbon atoms, and so on.

5 As used herein with respect to a substituting radical, and unless otherwise stated, the terms " C₂₋₇ alkenyl " and " aliphatic unsaturated hydrocarbon radical with 2 to 7 carbon atoms " are interchangeable and designate a straight and branched acyclic hydrocarbon monovalent radical having one or more ethylenical unsaturations and having from 2 to 7 carbon
10 atoms such as, for example, vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl, 2-hexenyl, 2-heptenyl, butadienyl, pentadienyl, hexadienyl, heptadienyl, heptatrienyl and the like, including all possible isomers thereof; the term " C₃₋₇ alkenyl " designate the corresponding radical with only 3 to 7 carbon atoms, and so on.

15 As used herein with respect to a substituting radical, and unless otherwise stated, the terms " C₃₋₁₀ cycloalkenyl" and " cycloaliphatic unsaturated hydrocarbon radical with 3 to 10 carbon atoms " are interchangeable and mean a monocyclic mono- or polyunsaturated hydrocarbon monovalent radical having from 3 to 8 carbon atoms, such as for
20 instance cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, cycloheptatrienyl, cyclooctenyl, cyclooctadienyl and the like, or a C₇₋₁₀ polycyclic mono- or polyunsaturated hydrocarbon mono-valent radical having from 7 to 10 carbon atoms such as dicyclopentadienyl, fencheryl (including all isomers
25 thereof, such as α -pinolenyl), bicyclo[2.2.1]hept-2-enyl, bicyclo[2.2.1]hepta-2,5-dienyl, cyclo-fencheryl and the like.

 As used herein with respect to a substituting radical, and unless otherwise stated, the term " C₂₋₇ alkynyl " defines straight and branched chain hydrocarbon radicals containing one or more triple bonds and having from 2 to
30 20 carbon atoms such as, for example, acetylenyl, 2-propynyl, 3-butyne, 2-butyne, 2-pentyne, 3-pentyne, 3-methyl-2-butyne, 3-hexynyl, 2-hexynyl and the like and all possible isomers thereof.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms "arylalkyl" and "heterocyclic-substituted alkyl" refer to an aliphatic saturated hydrocarbon monovalent radical, preferably a C₁₋₇ alkyl or a C₃₋₁₀ cycloalkyl such as defined above, onto which an aryl radical or respectively a heterocyclic radical (such as defined above) is already bonded, such as but not limited to benzyl, pyridylmethyl, pyridylethyl, 2-(2-pyridyl)isopropyl, oxazolylbutyl, 2-thienylmethyl and 2-furylmethyl.

As used herein with respect to a substituting radical, and unless otherwise stated, the term "alkylaryl" and "alkyl-substituted heterocyclic" refer to an aryl radical or respectively a heterocyclic radical (such as defined above) onto which is (are) already bonded one or more aliphatic saturated hydrocarbon monovalent radicals, preferably C₁₋₇ alkyl radicals or C₃₋₁₀ cycloalkyl radicals as defined above such as, but not limited to, o-toluy, m-toluy, p-toluy, mesityl and 2,4,6-trimethylphenyl.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms "alkylamino", "cycloalkylamino", "alkenyl-amino", "cycloalkenylamino", "arylamino", "arylalkylamino", "heterocyclic amino", "hydroxyalkylamino", "mercaptoalkylamino" and "alkynylamino" mean that respectively one (thus monosubstituted amino) or even two (thus disubstituted amino) C₁₋₇ alkyl, C₃₋₁₀ cycloalkyl, C₂₋₇ alkenyl, C₃₋₁₀ cycloalkenyl, aryl, arylalkyl, heterocyclic, mono- or polyhydroxy C₁₋₇ alkyl, mono- or polymercapto C₁₋₇ alkyl or C₂₋₇ alkynyl radical(s) (each of them as defined herein, respectively) is/are attached to a nitrogen atom through a single bond or, in the case of heterocyclic, include a nitrogen atom, such as but not limited to, anilino, benzylamino, methylamino, dimethylamino, ethylamino, diethylamino, isopropylamino, propenylamino, n-butylamino, ter-butylamino, dibutylamino, morpholino-alkylamino, morpholinyl, piperidiny, piperazinyl, hydroxymethylamino, β -hydroxyethylamino and ethynylamino; this definition also includes mixed disubstituted amino radicals wherein the nitrogen atom is attached to two such radicals belonging to two different sub-set of radicals, e.g. an alkyl radical and an alkenyl radical, or to two different radicals within the same sub-set of radicals, e.g. methylethylamino; the term "C₃₋₇ alkyl-amino" designates the corresponding radical with only 3 to 7 carbon atoms in

the alkyl group(s) attached to nitrogen, for instance di-isopropylamino, and so on; among disubstituted amino radicals, symmetrically substituted are usually preferred and more easily accessible.

As used herein with respect to a substituting radical, and unless otherwise stated, the terms "(thio)carboxylic acid ester", "(thio)carboxylic acid thioester" and "(thio)carboxylic acid amide" refer to radicals wherein the carboxyl or thiocarboxyl group is directly attached to the pteridine ring (e.g. in the 6- and/or 7-position) and wherein said carboxyl or thiocarboxyl group is bonded to the hydrocarbonyl residue of an alcohol, a thiol, a polyol, a phenol, a thiophenol, a primary or secondary amine, a polyamine, an amino-alcohol or ammonia, the said hydrocarbonyl residue being selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, alkylaryl, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, arylamino, arylalkylamino, heterocyclic amino, hydroxyalkylamino, mercapto-alkylamino or alkynylamino (such as above defined, respectively).

As used herein with respect to a substituting radical, and unless otherwise stated, the term "amino-acid" refers to a radical derived from a molecule having the chemical formula $H_2N-CHR-COOH$, wherein R is the side group of atoms characterizing the amino-acid type; said molecule may be one of the 20 naturally-occurring amino-acids or any similar non naturally-occurring amino-acid.

As used herein and unless otherwise stated, the term "stereoisomer" refers to all possible different isomeric as well as conformational forms which the compounds of formula (I) or (II) may possess, in particular all possible stereochemically and conformationally isomeric forms, all diastereomers, enantiomers and/or conformers of the basic molecular structure. Some compounds of the present invention may exist in different tautomeric forms, all of the latter being included within the scope of the present invention.

As used herein and unless otherwise stated, the term "enantiomer" means each individual optically active form of a compound of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (i.e. at least 90% of one enantiomer and at

most 10% of the other enantiomer), preferably at least 90% and more preferably at least 98%.

As used herein and unless otherwise stated, the term " solvate " includes any combination which may be formed by a pteridine derivative of
5 this invention with a suitable inorganic solvent (e.g. hydrates) or organic solvent, such as but not limited to alcohols, ketones, esters and the like.

As used herein and unless otherwise stated, the terms " dihydro- pteridine derivative " and " tetrahydropteridine derivative " refer to the hydrogenation products of the pteridine derivatives having formula (I) or (II),
10 i.e. derivatives wherein two hydrogen atoms are present in positions 5 and 6, or 7 and 8, of the pteridine ring, or wherein four hydrogen atoms are present in positions 5, 6, 7 and 8 of the said ring; such hydrogenated derivatives are easily accessible from the pteridine derivatives using hydrogenation methods well known in the art.

15

DETAILED DESCRIPTION OF THE INVENTION

A main object of the invention is to provide a pharmaceutical composition having high immunosuppressive activity. Thus, the present invention relates in particular to the medical applications of a group of pteridine derivatives, their
20 pharmaceutically acceptable salts, N-oxides, solvates, dihydro- and tetrahydroderivatives and enantiomers, possessing unexpectedly desirable pharmaceutical properties, in particular which are highly active immunosuppressive agents, and as such are useful in the treatment in transplant rejection and/or in the treatment of certain inflammatory diseases.

25 Surprisingly, the compounds of the present invention show a broader therapeutic spectrum profile than merely immunosuppressive activity, as is evidenced by the results obtained in the diversity of test procedures disclosed hereinbelow. A further advantageous feature of the compounds of the present invention resides in their excellent oral activity.

30 In the first embodiment of the invention, the novel pteridine derivatives are as defined in the general formula (I), wherein each of the substituents X, Z, R₁, R₂, R₃ and R₄ may correspond to any of the definitions given above (and, when X includes sulfur, wherein m may be 0, 1 or 2), in particular with

any of the individual meanings (such as illustrated above) of generic terms such as, but not limited to, "C₁₋₇ alkyl", "C₂₋₇ alkenyl", "C₂₋₇ alkynyl", "aryl", "alkylaryl", "arylalkyl", "alkylamino", "cycloalkylamino", "alkenylamino", "alkynylamino", "arylamino", "arylalkylamino", "C₁₋₇ alkoxy", "C₃₋₁₀ cycloalkoxy", "thio C₁₋₇ alkyl", "thio C₃₋₁₀ cycloalkyl", "halo C₁₋₇ alkyl", "amino-acid" and the like.

When a mixture of enantiomers of a pteridine derivative having the general formula (I) according to the invention is obtained during synthesis, the said mixture may be separated by means and methods standard in the art, e.g. liquid chromatography using one or more suitable chiral stationary phases. The latter include, for example, polysaccharides, in particular cellulose or amylose derivatives. Commercially available polysaccharide-based chiral stationary phases suitable for this purpose are ChiralCel™ CA, OA, OB, OC, OD, OF, OG, OJ and OK, and Chiralpak™ AD, AS, OP(+) and OT(+). Appropriate eluents or mobile phases for use in combination with said polysaccharide-based chiral stationary phases are hydrocarbons such as hexane and the like, optionally admixed with an alcohol such as ethanol, isopropanol and the like. The above mixture of enantiomers may alternatively be separated by forming diastereoisomers, followed by separation of the diastereoisomers, e.g. by differential crystallization or chromatography. The resolving agent may be cleaved from the separated diastereoisomers, e.g. by treatment with acids or bases, in order to generate the pure enantiomers of the compounds of the invention.

Some preferred pteridine derivatives having the general formula (I) or (II) according to the invention are more specifically illustrated in the following examples and defined in the following claims. For instance, useful pteridine species disclosed below include those wherein:

- R₁ is selected from the group consisting of methyl, ethyl, isopropyl, pentyl and benzyl, and/or
- R₂ is amino, and/or
- R₄ is hydrogen or methoxy, and/or
- R₃ is 3-thienyl or a phenyl group with one or more substituents (in the latter case, such substituents are preferably each independently selected from

the group consisting of fluoro, methoxy, ethoxy, trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate, and/or

- X is a sulfur atom (i.e. m is 0) or an oxygen atom, or
- 5 - X is NZ, wherein Z is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl and benzyl, or NZ together with R₁ forms a radical selected from the group consisting of hydroxylamino, morpholinyl, piperidinyl, piperazinyl, N-methylpiperazinyl, 1,2,4-triazolyl and pyrrolidinyl.

The present invention further provides processes and methods for making
 10 the novel pteridine derivatives having the general formula (I). As a general rule, the preparation of these compounds is based on the principle that, starting from a suitable pteridine precursor, each of the substituents XR₁, R₂, R₃ and R₄ may be introduced separately (except, of course, when R₃ together with R₄ forms a homocyclic or heterocyclic radical) without adversely
 15 influencing the presence of one or more substituents already introduced at other positions on the pteridine ring or the capacity to introduce further substituents later on.

Methods of manufacture have been developed by the present inventors which may be used alternatively to, or may be combined with, the methods of
 20 synthesis already known in the art of pteridine derivatives (depending upon the targeted final compound). For instance, methods for simultaneously introducing R₃ and R₄ in the form of a homocyclic or heterocyclic radical at positions 6 and 7 of the pteridine ring are already known from U.S. Patent No. 2,581,889. The synthesis of mono- and di-N-oxides of the pteridine derivatives
 25 of this invention can easily be achieved by treating the said derivatives with an oxidizing agent such as, but not limited to, hydrogen peroxide (e.g. in the presence of acetic acid) or a peracid such as chloroperbenzoic acid. Dihydro- and tetrahydropteridine derivatives of this invention can easily be obtained by catalytic hydrogenation of the corresponding pteridine derivatives, e.g. by
 30 placing the latter in a hydrogen atmosphere in the presence of platinum oxide or platinum. The methods for making the pteridine derivatives of the present invention will now be explained in more details by reference to the appended figures 1 to 5 wherein, unless otherwise stated hereinafter, each of the

substituting groups or atoms X, Z, R₁, R₂, R₃ and R₄ is as defined in formula (I) of the summary of the invention and, more specifically, may correspond to any of the individual meanings disclosed above. The same manufacturing methods may also be applied, if need be, while starting from pteridine derivatives which are already known in the art. In the description of the reaction steps involved in each figure, reference is made to the use of certain catalysts and/or certain types of solvents. It should be understood that each catalyst mentioned should be used in a catalytic amount well known to the skilled person with respect to the type of reaction involved. Solvents that may be used in the following reaction steps include various kinds of organic solvents such as protic solvents, polar aprotic solvents and non-polar solvents as well as aqueous solvents which are inert under the relevant reaction conditions. More specific examples include aromatic hydrocarbons, chlorinated hydrocarbons, ethers, aliphatic hydrocarbons, alcohols, esters, ketones, amides, water or mixtures thereof, as well as supercritical solvents such as carbon dioxide (while performing the reaction under supercritical conditions). The suitable reaction temperature and pressure conditions applicable to each kind of reaction step will not be detailed herein but do not depart from the relevant conditions already known to the skilled person with respect to the type of reaction involved and the type of solvent used (in particular its boiling point).

Figure 1 represents a scheme for the preparation of 2,4,6-trisubstituted pteridines with various R₂ and R₃ substituents in the 2- and 6-positions of the pteridine ring. In the first step (a), a chloropyrimidine 1, wherein R₂ may be *inter alia* amino, alkylamino, arylamino, alkoxy, aryloxy, mercaptoalkyl, or mercaptoaryl, is reacted with an appropriate nucleophile R₁XH, the said nucleophile being selected from the group consisting of alcohols (e.g. methanol, ethanol, isopropanol or benzylalcohol), thiols, primary amines and secondary amines wherein R₁ may be *inter alia* alkyl, cycloalkyl, aryl, alkylaryl, heteroaryl or alkylheteroaryl. Introduction of a nitroso group into the pyrimidine intermediate 2 occurs in step (b) under acidic aqueous conditions in the presence of sodium nitrite NaNO₂. Reduction of the nitroso functionality of the pyrimidine intermediate 3 into a free amino group in intermediate 4 is then

effected in step (c) by means of reducing agents (such as $\text{Na}_2\text{S}_2\text{O}_4$ or $(\text{NH}_4)_2\text{S}$) in water, or catalytically (Pt/H_2) in the presence of a protic solvent. In step (d), ring closure is performed by treating the diaminopyrimidine **4** with glyoxal in order to form a pteridine ring. In step (e), the nitrogen atom at
 5 position 8 of the pteridine ring of compound **5** is oxidized, e.g. using H_2O_2 under acidic conditions. In step (f), a chlorine atom is regioselectively introduced on the 6 position of the pteridine ring of compound **6** by treatment with a carboxylic acid chloride such as acetyl chloride under acidic conditions. Then in step (g) the 6-chlorosubstituted pteridine **7** is reacted with a boronic
 10 acid having the general formula $\text{R}_3\text{B}(\text{OH})_2$, wherein R_3 may be alkyl, cycloalkyl, aryl or heteroaryl, under basic conditions (such as in the presence of an aqueous alkaline solution) and a palladium based catalyst, thus yielding the desired derivative **8** of the present invention.

Figure 2 represents a scheme for the preparation of 2,4,6-trisubstituted
 15 pteridines with various R_2 and R_3 substituents in the 2- and 6-positions of the pteridine ring. In step (a), a nitroso group is introduced in position 5 of the pyrimidine ring of a compound **9**, wherein R_2 may be *inter alia* amino, alkylamino or arylamino, using sodium nitrite under aqueous acidic conditions. Reduction of the nitroso group of compound **10** in step (b) is achieved either
 20 catalytically (Pt/H_2) in the presence of a protic solvent, or chemically using sodium dithionite or ammonium sulfide in the presence of water. Then in the next step (c), the condensation of the diaminopyrimidine **11** with an α -ketoaldoxime bearing the group R_3 , wherein R_3 may be alkyl, cycloalkyl, aryl or heteroaryl, under acidic conditions in the presence of a protic solvent such
 25 as methanol, regioselectively yields the 6-substituted pteridine derivative **12**. Activation of the hydroxyl group of the tautomeric form of **12** by a nucleophilic displacement reaction occurs by preparing the 4-[(1,2,4)-triazolyl] pteridine derivative **13**, using POCl_3 or 4-chlorophenyl phosphorodichloridate, and 1,2,4-triazole in the presence of e.g. pyridine as solvent. If R_2 is a free amino
 30 group, protection of R_2 e.g. by means of an acetyl group may be necessary before carrying out the reaction of step (d), followed by a deprotection of the acetyl group during the nucleophilic displacement reaction. The nucleophilic substitution in step (e) may be performed, e.g. in the presence of 1,4-dioxane

as a solvent, by mixing the pteridine derivative **13** at room temperature with an appropriate nucleophile R_1XH , the said nucleophile being selected from the group consisting of alcohols (e.g. methanol, ethanol, isopropanol or benzylalcohol), thiols, primary amines and secondary amines wherein R_1 may
 5 be *inter alia* alkyl, cycloalkyl, aryl, alkylaryl, heteroaryl or alkylheteroaryl, thus yielding the desired derivative **14** of the present invention.

Figure 3 represents a scheme for the preparation of 2,4,7-trisubstituted pteridines with various R_2 and R_4 substituents in the 2- and 7-positions of the pteridine ring. In reaction step (a), a tetraaminopyrimidine **15**, wherein R_2 may
 10 be *inter alia* amino, alkylamino or arylamino, is reacted with *inter alia* an alkylglyoxal, arylglyoxal, alkylarylglyoxal, heteroarylglyoxal or alkyl heteroarylglyoxal in water under basic conditions, yielding the 7-substituted pteridine derivative **16** wherein R_4 may be *inter alia* alkyl, cycloalkyl, aryl, heteroaryl, alkylaryl or alkylheteroaryl. In step (b), hydrolysis under reflux in
 15 the presence of sodium hydroxide 1N yielded the 4-hydroxy tautomer of pteridine derivative **17**. In step (c), a 1,2,4-triazolyl group is introduced in position 4 by reacting **17** with triazole in the presence of 4-chlorophenyl phosphorodichloridate and dry pyridine. Finally in reaction step (d), the 1,2,4-triazolyl group at position 4 of the pteridine derivative **18** is displaced by an
 20 appropriate nucleophile R_1XH , the said nucleophile being selected from the group consisting of alcohols (e.g. methanol, ethanol, isopropanol or benzylalcohol), thiols, primary amines and secondary amines wherein R_1 may be *inter alia* alkyl, cycloalkyl, aryl, alkylaryl, heteroaryl or alkylheteroaryl, in the presence of a polar aprotic or protic solvent, thus yielding the desired
 25 derivative **19** of the present invention.

Figure 4 represents a scheme for the synthesis of unsymmetrical 2,4,6-trisubstituted and 2,4,7-trisubstituted pteridine derivatives with various R_2 , R_3 and R_4 substituents in the 2-, 6- and 7-positions of the pteridine ring. In step (a), the thiol function on 2-mercapto-4,6-diaminopyrimidine is methylated, e.g.
 30 by reaction with methyl iodide in the presence of a solvent such as ethanol, in order to yield 2-thiomethyl-4,6-diaminopyrimidine. Introduction of a nitroso group in the 5-position of the pyrimidine ring is then achieved in step (b) by using sodium nitrite under aqueous acidic conditions. In the next step (c), the

methylthio group in the 2-position is exchanged for a group R_2 , wherein R_2 may be *inter alia* alkoxy, aryloxy, cycloalkyloxy, heteroaryloxy, mercaptoalkyl, mercaptoaryl, mercaptocyclo-alkyl or mercaptoheteroaryl by reaction with an appropriate nucleophile. Reduction of the nitroso functionality is then achieved
 5 in step (d) either catalytically (Pt/H_2) in the presence of a protic solvent or chemically using sodium dithionite or ammonium sulfide in the presence of water. Then in the next step (e), the condensation of the 2-substituted-4,5,6-triaminopyrimidine with an α -ketoaldoxime bearing the group R_3 , wherein R_3 may be *inter alia* alkyl, cycloalkyl, aryl or heteroaryl, under acidic conditions in
 10 the presence of a solvent such as methanol, regioselectively yields a 2,6-substituted-4-aminopteridine derivative. The corresponding 2,7-substituted-4-aminopteridine can be obtained, according to step (i), by reaction of the 2-substituted-4,5,6-triaminopyrimidine with a glyoxal bearing the group R_4 , wherein R_4 may be *inter alia* alkyl, cycloalkyl, aryl or heteroaryl. According to
 15 step (f), acidic or basic hydrolysis of the amino group at position 4 of the pteridine ring, performed on the derivative from step (e) or (i), yields the corresponding 4-oxopteridine derivative. In step (g), the hydroxyl group of the tautomeric form of the latter is activated by nucleophilic displacement, e.g. by preparing the 4-[(1,2,4)-triazolyl] pteridine derivative. Finally, the nucleophilic
 20 displacement in step (h) is performed by mixing the said 4-triazolyl pteridine derivative with the appropriate nucleophile R_1XH , wherein R_1 may be *inter alia* alkyl, cycloalkyl, aryl, heteroaryl, alkylaryl, alkylheteroaryl.

Figure 5 represents a scheme for the synthesis of symmetrical 2,4,6-trisubstituted and 2,4,7-trisubstituted pteridine derivatives with various R_2 , R_3
 25 and R_4 substituents in the 2-, 6- and 7-positions of the pteridine ring. In step (a), the pyrimidine ring is nitrated in position 5 under strongly acidic conditions (HNO_3 ; H_2SO_4). Then in step (b) both hydroxyl groups (from the tautomeric form) are converted to chloro groups by treatment with a chlorinating agent such as $POCl_3$ or $SOCl_2$. Both chloro substituents are then displaced in step
 30 (c) with an appropriate nucleophile R_1XH , wherein R_1 may be *inter alia* alkyl, cycloalkyl, aryl, heteroaryl, alkylaryl, alkylheteroaryl. The nitro group is reduced in step (d) to an amino group by treatment with a reducing agent (e.g. Pt/H_2). Finally, reaction of the 2,4-substituted-5,6-diaminopyrimidine from step

(d) with an α -ketoaldoxime bearing the group R_3 , wherein R_3 may be *inter alia* alkyl, cycloalkyl, aryl or heteroaryl, regioselectively yields the desired 2,4,6-trisubstituted pteridine derivative in step (e). Alternatively, reaction of the 2,4-substituted-5,6-diaminopyrimidine from step (d) with a glyoxal bearing the
 5 group R_4 , wherein R_4 may be *inter alia* alkyl, cycloalkyl, aryl or heteroaryl, yields the desired 2,4,7-trisubstituted pteridine derivative in step (f).

Some sub-sets of pteridine derivatives according to the invention deserve specific interest. Thus in a particular embodiment the invention relates to a group of pteridine derivatives, as well as pharmaceutical
 10 compositions comprising such pteridine derivatives as active principle, having the above general formula (I) wherein:

- X is NZ,
- XR_1 is selected from the group consisting of hydroxylamino; hydrazino (i.e. Z is hydrogen and R_1 is amino); (mono- or di-) C_{1-7} alkylamino (i.e. R_1 is
 15 C_{1-7} alkyl); (mono- or di-) arylamino (i.e. R_1 is aryl, e.g. adamantyl); (mono- or di-) C_{3-10} cycloalkylamino (i.e. R_1 is C_{3-10} cycloalkyl); (mono- or di-) hydroxy C_{1-7} alkylamino (i.e. R_1 is C_{1-7} alkyl substituted with hydroxyl); (mono- or di-) C_{1-4} alkylarylamino; and saturated or unsaturated heterocyclic groups containing at least one nitrogen atom and optionally
 20 substituted by one or more substituents independently selected from the group consisting of C_{1-4} alkyl, hydroxy C_{1-4} alkyl, C_{1-4} alkyloxy, halogen, hydroxy, hydroxycarbonyl and C_{1-4} alkyloxycarbonyl, such as but not limited to piperazinyl, N-alkylpiperazinyl and morpholinyl;
- R_2 is amino or is selected from the group consisting of hydroxylamino; hydrazino (i.e. Z is hydrogen and R_1 is amino); (mono- or di-) arylamino
 25 (i.e. R_1 is aryl); (mono- or di-) C_{3-10} cycloalkylamino (i.e. R_1 is C_{3-10} cycloalkyl); (mono- or di-) hydroxy C_{1-7} alkylamino (i.e. R_1 is C_{1-7} alkyl substituted with hydroxyl); (mono- or di-) C_{1-4} alkylarylamino; and saturated or unsaturated heterocyclic groups containing at least one nitrogen atom
 30 and optionally substituted by one or more substituents independently selected from the group consisting of C_{1-4} alkyl, hydroxy C_{1-4} alkyl, C_{1-4} alkyloxy, halogen, hydroxy, hydroxycarbonyl and C_{1-4} alkyloxycarbonyl, such as but not limited to piperazinyl, N-alkylpiperazinyl and morpholinyl;

- R_3 is selected from the group consisting of unsubstituted, monosubstituted and disubstituted aryl groups (wherein the substituent(s) may independently be halogen, C_{1-4} alkoxy or C_{1-4} alkyl); 3,4,5-trimethoxyphenyl; 3,4-formylidene-3,4-dihydroxyphenyl; aryl groups bonded to the pteridine ring via a saturated or unsaturated aliphatic spacer which may be halogenated or hydroxylated; and aliphatic substituents which may contain ether function, alcohol function, or substituted or unsubstituted amino functions or C_{1-4} alkyloxy; and
 - R_4 is selected from the group consisting of hydrogen, alkyl and alkoxy;
- 10 or a pharmaceutically acceptable addition salt or a stereochemical isomeric form thereof.

In another particular embodiment, the invention relates to a group of pteridine derivatives, as well as pharmaceutical compositions comprising such pteridine derivatives as active principle, having the above general formula (I) wherein:

- X is oxygen or sulfur;
- R_1 is C_{1-7} alkyl;
- R_2 is amino or is selected from the group consisting of hydroxylamino; hydrazino (i.e. Z is hydrogen and R_1 is amino); (mono- or di-) arylamino (i.e. R_1 is aryl); (mono- or di-) C_{3-10} cycloalkylamino (i.e. R_1 is C_{3-10} cycloalkyl); (mono- or di-) hydroxy C_{1-7} alkylamino (i.e. R_1 is C_{1-7} alkyl substituted with hydroxyl); (mono- or di-) C_{1-4} alkylarylamino; and saturated or unsaturated heterocyclic groups containing at least one nitrogen atom and optionally substituted by one or more substituents independently selected from the group consisting of C_{1-4} alkyl, hydroxy C_{1-4} alkyl, C_{1-4} alkyloxy, halogen, hydroxy, hydroxycarbonyl and C_{1-4} alkyloxycarbonyl, such as but not limited to piperazinyl, N-alkylpiperazinyl and morpholinyl;
- R_3 is selected from the group consisting of unsubstituted, monosubstituted and disubstituted aryl groups (wherein the substituent(s) may independently be halogen, C_{1-4} alkoxy or C_{1-4} alkyl); aryl groups bonded to the pteridine ring via a saturated or unsaturated aliphatic spacer which may be halogenated or hydroxylated; and aliphatic substituents which may contain

ether function, alcohol function, or substituted or unsubstituted amino functions or C₁₋₄ alkyloxy; and

- R₄ is selected from the group consisting of hydrogen, alkyl and alkoxy; or a pharmaceutically acceptable addition salt or a stereochemical isomeric form thereof.

In another particular embodiment, the invention relates to a group of pteridine derivatives having the above general formula (I) wherein X represents an oxygen atom or a group with the formula S(O)_m, wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R₁ is a group selected from the group consisting of C₁₋₄ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₄ alkyl, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkyl-amino, hydroxylalkylamino, mercaptoalkyl-amino, heterocyclic amino, hydrazino, alkylhydrazino and phenyl-hydrazino; or R₁ is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;
- Z is a group independently defined as R₁ or Z is hydrogen or the group NZ together with R₁ is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R₂ is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl; ureido; thio-ureido; sulfonamido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino;

- alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C₃₋₇ alkylamino; arylamino; arylalkyl-amino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; thio C₁₋₇ alkyl; 5 arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C₃₋₁₀ cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals;
- R₄ is an atom or a group selected from the group consisting of hydrogen; halogen; C₁₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; carboxy C₁₋₇ 10 alkyl; carboxyaryl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₁₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-acetal; carboxylic acid; carboxylic 15 acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; 20 hydroxyalkylamino; mercaptoalkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkylloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and thioamides thereof, aryl being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₇ 25 alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, 30 hydroxylamino, alkoxyamino, mercaptoamino, thioalkylamino, acylamino, thioacyl-amino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino,

alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino,
 hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino,
 alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic
 radicals; aromatic or heterocyclic substituents substituted with an aliphatic
 5 spacer between the pteridine ring and the aromatic or heterocyclic
 substituent, whereby said aliphatic spacer is a branched or straight,
 saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may
 contain one or more functions, atoms or radicals selected from the group
 consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether,
 10 thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-
 acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido,
 thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or
 amide, thiocarboxylic acid or ester or thioester or halide or anhydride or
 amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino,
 15 mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenyl-
 amino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino,
 mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino,
 phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or
 straight, saturated or unsaturated aliphatic chains of 1 to 7 carbon atoms
 20 optionally containing one or more functions selected from the group
 consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether,
 thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino,
 aminoacid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl,
 ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide,
 25 thiocarboxylic acid or ester or thioester or halide or anhydride or amide,
 nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino,
 alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynyl-
 amino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino,
 heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl,
 30 sulfinyl, sulfonamido and halogen; and
 - R₃ is selected from chloro and thiomethyl;

and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-*N*-oxide thereof and/or a solvate and/or a dihydro- or tetrahydropteridine derivative thereof.

When applicable, and depending upon the specific substituents being present, not only the novel pteridine having the general formula (I) but also some pteridine derivatives previously known in the art without any indication of biological activity, i.e. all of the pteridines having the general formula (II) according to this invention, may be in the form of a pharmaceutically acceptable salt. The latter include any therapeutically active non-toxic addition salt which compounds having the general formula (II) are able to form with a salt-forming agent. Such addition salts may conveniently be obtained by treating the pteridine derivatives of the invention with an appropriate salt-forming acid or base. For instance, pteridine derivatives having basic properties may be converted into the corresponding therapeutically active, non-toxic acid addition salt form by treating the free base form with a suitable amount of an appropriate acid following conventional procedures. Examples of such appropriate salt-forming acids include, for instance, inorganic acids resulting in forming salts such as but not limited to hydrohalides (e.g. hydrochloride and hydrobromide), sulfate, nitrate, phosphate, diphosphate, carbonate, bicarbonate, and the like; and organic monocarboxylic or dicarboxylic acids resulting in forming salts such as, for example, acetate, propanoate, hydroxyacetate, 2-hydroxypropanoate, 2-oxopropanoate, lactate, pyruvate, oxalate, malonate, succinate, maleate, fumarate, malate, tartrate, citrate, methanesulfonate, ethanesulfonate, benzoate, 2-hydroxybenzoate, 4-amino-2-hydroxybenzoate, benzene-sulfonate, *p*-toluenesulfonate, salicylate, *p*-aminosalicylate, pamoate, bitartrate, camphorsulfonate, edetate, 1,2-ethanedisulfonate, fumarate, glucoheptonate, gluconate, glutamate, hexylresorcinate, hydroxynaphtoate, hydroxyethanesulfonate, mandelate, methylsulfate, pantothenate, stearate, as well as salts derived from ethanedioic, propanedioic, butanedioic, (*Z*)-2-butenedioic, (*E*)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic and cyclohexanesulfamic acids and the like.

Pteridine derivatives having acidic properties may be converted in a similar manner into the corresponding therapeutically active, non-toxic base addition salt form. Examples of appropriate salt-forming bases include, for instance, inorganic bases like metallic hydroxides such as but not limited to those of alkali and alkaline-earth metals like calcium, lithium, magnesium, potassium and sodium, or zinc, resulting in the corresponding metal salt; organic bases such as but not limited to ammonia, alkylamines, benzathine, hydrabamine, arginine, lysine, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylene-diamine, N-methylglucamine, procaine and the like.

Reaction conditions for treating the pteridine derivatives (II) of this invention with an appropriate salt-forming acid or base are similar to standard conditions involving the same acid or base but different organic compounds with basic or acidic properties, respectively. Preferably, in view of its use in a pharmaceutical composition or in the manufacture of medicament for treating specific diseases, the pharmaceutically acceptable salt will be designed, i.e. the salt-forming acid or base will be selected so as to impart greater water-solubility, lower toxicity, greater stability and/or slower dissolution rate to the pteridine derivative of this invention.

The present invention further provides the use of a pteridine derivative represented by the general formula (II), or a pharmaceutically acceptable salt or a solvate thereof, as a biologically-active ingredient, i.e. an active principle, especially as a medicine or a diagnostic agent or for the manufacture of a medicament or a diagnostic kit. In particular the said medicament may be for the prevention or treatment of a pathologic condition selected from the group consisting of:

- immune disorders, in particular organ and cells transplant rejections, and autoimmune disorders,
- cardiovascular disorders,
- allergic conditions,
- disorders of the central nervous system, and
- cell proliferative disorders.

The pathologic conditions and disorders concerned by the said use, and the corresponding methods of prevention or treatment, are detailed hereinbelow. Any of the uses mentioned with respect to the present invention may be restricted to a non-medical use (e.g. in a cosmetic composition), a
 5 non-therapeutic use, a non-diagnostic use, a non-human use (e.g. in a veterinary composition), or exclusively an *in-vitro* use, or a use with cells remote from an animal.

The invention further relates to a pharmaceutical composition comprising:

- (a) one or more pteridine represented by the general formula (II), and
- 10 (b) one or more pharmaceutically acceptable carriers.

In a third embodiment, this invention provides combinations, preferably synergistic combinations, of one or more pteridine represented by the general formula (II) with one or more biologically-active drugs being preferably selected from the group consisting of immunosuppressant and/or
 15 immunomodulator drugs, antineoplastic drugs, anti-histamines, inhibitors of allergy-causative agents (anti-allergic drugs) and antiviral agents. As is conventional in the art, the evaluation of a synergistic effect in a drug combination may be made by analyzing the quantification of the interactions between individual drugs, using the median effect principle described by Chou
 20 et al. in *Adv. Enzyme Reg.* (1984) 22:27. Briefly, this principle states that interactions (synergism, additivity, antagonism) between two drugs can be quantified using the combination index (hereinafter referred as CI) defined by the following equation:

$$CI_x = \frac{ED_x^{1c}}{ED_x^{1a}} + \frac{ED_x^{2c}}{ED_x^{2a}}$$

25 wherein ED_x is the dose of the first or respectively second drug used alone (1a, 2a), or in combination with the second or respectively first drug (1c, 2c), which is needed to produce a given effect. The said first and second drug have synergistic or additive or antagonistic effects depending upon $CI < 1$, $CI = 1$, or $CI > 1$, respectively. As will be explained in more detail herein-below,
 30 this principle may be applied to a number of desirable effects such as, but not limited to, an activity against transplant rejection, an activity against

immunosuppression or immunomodulation, an activity against allergy or an activity against cell proliferation.

For instance the present invention relates to a pharmaceutical composition or combined preparation having synergistic effects against immuno-
5 suppression or immunomodulation and containing:

- (a) one or more immunosuppressant and/or immunomodulator drugs, and
- (b) at least one pteridine derivative represented by the general formula (II),
and
- (c) optionally one or more pharmaceutical excipients or pharmaceutically
10 acceptable carriers,

for simultaneous, separate or sequential use in the treatment or prevention of autoimmune disorders and/or in transplant-rejections.

Suitable immunosuppressant drugs for inclusion in the synergistic compositions or combined preparations of this invention belong to a well
15 known therapeutic class. They are preferably selected from the group consisting of cyclosporin A, substituted xanthines (e.g. methylxanthines such as pentoxifylline), daltroban, sirolimus, tacrolimus, rapamycin (and derivatives thereof such as defined below), leflunomide (or its main active metabolite A771726, or analogs thereof called malononitrilamides), mycophenolic acid
20 and salts thereof (including the sodium salt marketed under the trade name Mofetil®), adrenocortical steroids, azathioprine, brequinar, gusperimus, 6-mercaptopurine, mizoribine, chloroquine, hydroxychloroquine and monoclonal antibodies with immunosuppressive properties (e.g. etanercept, infliximab or kineret). Adrenocortical steroids within the meaning of this invention mainly
25 include glucocorticoids such as but not limited to ciprocinonide, desoxycorticosterone, fludrocortisone, flumoxonide, hydrocortisone, naflocort, procinonide, timobesone, tipredane, dexamethasone, methylprednisolone, methotrexate, prednisone, prednisolone, triamcinolone and pharmaceutically acceptable salts thereof. Rapamycin derivatives as referred herein include O-
30 alkylated derivatives, particularly 9-deoxorapamycins, 26-dihydrorapamycins, 40-O-substituted rapamycins and 28,40-O,O-disubstituted rapamycins (as disclosed in U.S. Patent No. 5,665,772) such as 40-O-(2-hydroxy) ethyl rapamycin – also known as SDZ-RAD -, pegylated rapamycin (as disclosed in

U.S. Patent No. 5,780,462), ethers of 7-desmethyrapamycin (as disclosed in U.S. Patent No. 6,440,991) and polyethylene glycol esters of SDZ-RAD (as disclosed in U.S. Patent No. 6,331,547).

Suitable immunomodulator drugs for inclusion into the synergistic immunomodulating pharmaceutical compositions or combined preparations of this invention are preferably selected from the group consisting of acemannan, amiprilose, bucillamine, dimepranol, ditiocarb sodium, imiquimod, Inosine Pranobex, interferon- β , interferon- γ , lentinan, levamisole, lisophylline, pidotimod, romurtide, platonin, procodazole, propagermanium, thymomodulin, thymopentin and ubenimex.

Synergistic activity of the pharmaceutical compositions or combined preparations of this invention against immunosuppression or immunomodulation may be readily determined by means of one or more lymphocyte activation tests. Usually activation is measured via lymphocyte proliferation. Inhibition of proliferation thus always means immunosuppression under the experimental conditions applied. There exist different stimuli for lymphocyte activation, in particular:

- a) co-culture of lymphocytes of different species (mixed lymphocyte reaction, hereinafter referred as MLR) in a so-called mixed lymphocyte culture test: lymphocytes expressing different minor and major antigens of the HLA-DR type (= alloantigens) activate each other non-specifically;
- b) a CD3 assay wherein there is an activation of the T-lymphocytes via an exogenously added antibody (OKT3). This antibody reacts against a CD3 molecule located on the lymphocyte membrane which has a co-stimulatory function. Interaction between OKT3 and CD3 results in T-cell activation which proceeds via the Ca^{2+} /calmodulin/calcineurin system and can be inhibited e.g. by cyclosporin A (hereinafter referred as CyA);
- c) a CD28 assay wherein specific activation of the T-lymphocyte proceeds via an exogenously added antibody against a CD28 molecule which is also located on the lymphocyte membrane and delivers strong co-stimulatory signals. This activation is Ca^{2+} -independent and thus cannot be inhibited by CyA.

Determination of the immunosuppressing or immunomodulating activity of the pteridine derivatives of this invention, as well as synergistic combinations comprising them, is preferably based on the determination of one or more, preferably at least three lymphocyte activation *in vitro* tests, more preferably including at least one of the MLR test, CD3 assay and CD28 assay referred above. Preferably the lymphocyte activation *in vitro* tests used include at least two assays for two different clusters of differentiation preferably belonging to the same general type of such clusters and more preferably belonging to type I transmembrane proteins. Optionally the determination of the immunosuppressing or immunomodulating activity may be performed on the basis of other lymphocyte activation *in vitro* tests, for instance by performing a TNF- α assay or an IL-1 assay or an IL-6 assay or an IL-10 assay or an IL-12 assay or an assay for a cluster of differentiation belonging to a further general type of such clusters and more preferably belonging to type II transmembrane proteins such as, but not limited to, CD69, CD 71 or CD134.

The synergistic effect may be evaluated by the median effect analysis method described herein-before. Such tests may for instance, according to standard practice in the art, involve the use of equipment, such as flow cytometer, being able to separate and sort a number of cell subcategories at the end of the analysis, before these purified batches can be analysed further.

Synergistic activity of the pharmaceutical compositions of this invention in the prevention or treatment of transplant rejection may be readily determined by means of one or more leukocyte activation tests performed in a Whole Blood Assay (hereinafter referred as WBA) described for instance by Lin et al. in *Transplantation* (1997) 63:1734-1738. WBA used herein is a lymphoproliferation assay performed *in vitro* using lymphocytes present in the whole blood, taken from animals that were previously given the pteridine derivative, and optionally the other immunosuppressant drug, *in vivo* (more details are given in example 118). Hence this assay reflects the *in vivo* effect of substances as assessed by an *in vitro* read-out assay. The synergistic effect may be evaluated by the median effect analysis method described herein-before. Various organ transplantation models in animals are also available *in vivo*, which are strongly influenced by different immunogenicities,

depending on the donor and recipient species used and depending on the nature of the transplanted organ. The survival time of transplanted organs can thus be used to measure the suppression of the immune response.

The pharmaceutical composition or combined preparation with synergistic activity against immunosuppression or immunomodulation according to this invention may contain the pteridine derivative of formula (II) over a broad content range depending on the contemplated use and the expected effect of the preparation. Generally, the pteridine derivative content of the combined preparation is within the range of 0.1 to 99.9% by weight, preferably from 1 to 99% by weight, more preferably from 5 to 95% by weight.

The invention further relates to a composition or combined preparation having synergistic effects against cell proliferation and containing:

- (a) one or more antineoplastic drugs, and
- (b) at least one pteridine derivative represented by the general formula (II), and
- (c) optionally one or more pharmaceutical excipients or pharmaceutically acceptable carriers,

for simultaneous, separate or sequential use in the treatment or prevention of cell proliferative disorders.

Suitable antineoplastic drugs for inclusion into the synergistic antiproliferative pharmaceutical compositions or combined preparations of this invention are preferably selected from the group consisting of alkaloids, alkylating agents (including but not limited to alkyl sulfonates, aziridines, ethylenimines, methylmelamines, nitrogen mustards and nitrosoureas), antibiotics, antimetabolites (including but not limited to folic acid analogs, purine analogs and pyrimidine analogs), enzymes, interferon and platinum complexes. More specific examples include acivicin; aclarubicin; acodazole; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene; bisnafide; bizelesin; bleomycin; brequinar; bropiramine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin; carzelesin; cedefingol; chlorambucil;

cirolemycin; cisplatin; cladribine; crisnatol; cyclophosphamide; cytarabine;
 dacarbazine; dactinomycin; daunorubicin; decitabine; dexormaplatin;
 dezaguanine; diaziquone; docetaxel; doxorubicin; droloxifene;
 dromostanolone; duazomycin; edatrexate; eflomithine; elsamitrucin;
 5 enloplatin; enpromate; epiropidine; epirubicin; erbulozole; esorubicin;
 estramustine; etanidazole; ethiodized oil I 131; etoposide; etoprine; fadrozole;
 fazarabine; fenretinide; floxuridine; fludarabine; fluorouracil; flurocitabine;
 fosquidone; fostriecin; gemcitabine; Gold 198; hydroxyurea; idarubicin;
 ifosfamide; ilmofofine; interferon α -2a; interferon α -2b; interferon α -n1;
 10 interferon α -n3; interferon β -1a; interferon γ -1b; iproplatin; irinotecan;
 lanreotide; letrozole; leuprolide; liarozole; lometrexol; lomustine; losoxantrone;
 masoprocol; maytansine; mechlorethamine; megestrol; melengestrol;
 melphalan; menogaril; mercaptopurine; methotrexate; metoprine; meturedopa;
 mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin;
 15 mitosper; mitotane; mitoxantrone; mycophenolic acid; nocodazole; nogala-
 mycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin;
 pentamustine; peplomycin; perfosfamide; pipobroman; piposulfan;
 piroxantrone; plicamycin; plomestane; porfimer; porfiromycin; prednimustine;
 procarbazine; puromycin; pyrazofurin; riboprine; rogletimide; safingol;
 20 semustine; simtrazene; sparfosate; sparsomycin; spirogermanium;
 spiromustine; spiroplatin; streptonigrin; streptozocin; strontium 89 chloride;
 sulofenur; talisomycin; taxane; taxoid; tecogalan; tegafur; teloxantrone;
 temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine;
 thiotepa; tiazofurin; tirapazamine; topotecan; toremifene; trestolone; tricitabine;
 25 trimetrexate; triptorelin; tubulozole; uracil mustard; uredepa; vapreotide;
 verteporfin; vinblastine; vincristine; vindesine; vinepidine; vinglycinate;
 vinleurosine; vinorelbine; vinrosidine; vinzolidine; vorozole; zeniplatin;
 zinostatin; zorubicin; and their pharmaceutically acceptable salts.

Other suitable anti-neoplastic compounds include 20-epi-1,25
 30 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene;
 adecyphenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine;
 ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine;
 anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist

- D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; anti-androgens such as, but not limited to, benorterone, cioteronel, cyproterone, delmadinone, oxendolone, topterone, zanoterone and their pharmaceutically acceptable salts; anti-estrogens such as, but not limited to, clometherone;
- 5 delmadinone; nafoxidine; nitromifene; raloxifene; tamoxifen; toremifene; trioxifene and their pharmaceutically acceptable salts; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin; azasetron; azatoxin;
- 10 azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstaurosporine; β -lactam derivatives; β -alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; broprimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C;
- 15 camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors; castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; clomifene and analogues thereof; clotrimazole; collismycin A and B;
- 20 combretastatin and analogues thereof; conagenin; crambescidin 816; cryptophycin and derivatives thereof; curacin A; cyclopentantraquinones; cycloplatam; cypemycin; cytarabine; cytolytic factor; cytostatin; dacliximab; dehydroidemnin B; deslorelin; dexifosfamide; dexrazoxane; dexverapamil; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol;
- 25 dioxamycin; diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; elemene; emitefur; epristeride; estrogen agonists and antagonists; exemestane; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fluorodaunorunicin; forfenimex; formestane; fotemustine;
- 30 gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idoxifene; idramantone; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth

factor-1 receptor inhibitor; interferon agonists; iobenguane; iododoxorubicin; ipomeanol; irinotecan; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N; leinamycin; lenograstim; lentinan; leptolstatin; leukemia inhibiting factor; leuprorelin; 5 levamisole; liarozole; lissoclinamide; lobaplatin; lombricine; lonidamine; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; mannostatin A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitors; mifepristone; miltefosine; mirimostim; 10 mitoguazone; mitolactol; mitonafide; mitotoxin fibroblast growth factor-saporin; mofarotene; molgramostim; human chorionic gonadotrophin monoclonal antibody; mopidamol; mycaperoxide B; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone; pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral 15 endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; octreotide; okicenone; onapristone; ondansetron; ondansetron; oracin; osaterone; oxaliplatin; oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; peldesine; pentosan; pentostatin; pentrozole; perflubron; perillyl 20 alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine; pirarubicin; piritrexim; placetin A and B; plasminogen activator inhibitor; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein kinase C inhibitors; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; raltitrexed; ramosetron; 25 ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitors; retelliptine; rhenium 186 etidronate; rhizoxin; retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; saintopin; sarcophytol A; sargramostim; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; 30 splenopentin; spongistatin 1; squalamine; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; suradista; suramin; swainsonine; tallimustine; tamoxifen; tauromustine; tazarotene; tecogalan; tellurapyrylium; telomerase inhibitors; temozolomide; tetrachlorodecaoxide;

tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; titanocene; topsentin; tretinoin; triacetyluridine; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; ubenimex; urogenital sinus-
 5 derived growth inhibitory factor; urokinase receptor antagonists; variolin B; velaresol; veramine; verdins; verteporfin; vinxaltine; vitaxin; zanoterone; zilascorb; and their pharmaceutically acceptable salts.

Synergistic activity of the pharmaceutical compositions or combined preparations of this invention against cell proliferation may be readily
 10 determined by means of one or more tests such as, but not limited to, the measurement of the radioactivity resulting from the incorporation of ^3H -thymidine in culture of tumor cell lines. For instance, different tumor cell lines are selected in order to evaluate the anti-tumor effects of the test compounds, such as but not limited to:

- 15 - RPMI1788: human Peripheral Blood Leucocytes (PBL) Caucasian tumor line,
- Jurkat: human acute T cell leukemia,
- EL4: C57Bl/6 mouse lymphoma, or
- THP-1: human monocyte tumor line.

20 Depending on the selected tumor cell line, different culture media may be used, such as for example:

- for RPMI1788 and THP-1: RPMI-1640 + 10% FCS + 1% NEAA + 1% sodium pyruvate + 5×10^{-5} mercapto-ethanol + antibiotics (G-418 0.45 $\mu\text{g/ml}$).
- 25 - for Jurkat and EL4: RPMI-1640 + 10% FCS + antibiotics (G-418 0.45 $\mu\text{g/ml}$).

In a specific embodiment of the synergy determination test, the tumor cell lines are harvested and a suspension of 0.27×10^6 cells/ml in whole medium is prepared. The suspensions (150 μl) are added to a microtiter plate in triplicate.
 30 Either complete medium (controls) or the test compounds at the test concentrations (50 μl) are added to the cell suspension in the microtiter plate. The cells are incubated at 37°C under 5% CO_2 for about 16 hours. ^3H -

thymidine is added, and the cells incubated for another 8 hours. The cells are harvested and radioactivity is measured in counts per minute (CPM) in a β -counter. The ^3H -thymidine cell content, and thus the measured radioactivity, is proportional to the proliferation of the cell lines. The synergistic effect is evaluated by the median effect analysis method as disclosed herein-before.

The pharmaceutical composition or combined preparation with synergistic activity against cell proliferation according to this invention may contain the pteridine compound of formula (II) over a broad content range depending on the contemplated use and the expected effect of the preparation. Generally, the pteridine content of the combined preparation is within the range of 0.1 to 99.9% by weight, preferably from 1 to 99% by weight, more preferably from 5 to 95% by weight.

The invention further relates to a pharmaceutical composition or combined preparation having synergistic effects against a viral infection and containing:

- (a) one or more anti-viral agents, and
- (b) at least one pteridine derivative represented by the general formula (II), and
- (c) optionally one or more pharmaceutical excipients or pharmaceutically acceptable carriers,

for simultaneous, separate or sequential use in the treatment or prevention of a viral infection.

Suitable anti-viral agents for inclusion into the synergistic antiviral compositions or combined preparations of this invention include, for instance, retroviral enzyme inhibitors belonging to categories well known in the art, such as HIV-1 IN inhibitors, nucleoside reverse transcriptase inhibitors (e.g. zidovudine, lamivudine, didanosine, stavudine, zalcitabine and the like), non-nucleoside reverse transcriptase inhibitors (e.g. nevirapine, delavirdine and the like), other reverse transcriptase inhibitors (e.g. foscarnet sodium and the like), and HIV-1 protease inhibitors (e.g. saquinavir, ritonavir, indinavir, nelfinavir and the like). Other suitable antiviral agents include for instance acemannan, acyclovir, adefovir, alovudine, alvircept, amantadine, arantol, arildone, atevirdine, avaridine, cidofovir, cipamfylline, cytarabine, desciclovir, disoxanil, edoxudine, enviroxime, famciclovir, famotone,

fiacitabine, fialuridine, floxuridine, fosarilate, fosfonet, ganciclovir, idoxuridine, kethoxal, lobucavir, memotine, methisazone, penciclovir, pirodavid, somantadine, sorivudine, tilorone, trifluridine, valaciclovir, vidarabine, viroxime, zinviroxime, moroxydine, podophyllotoxin, ribavirine, rimantadine, 5 stallimycine, statolon, tromantadine and xenazoic acid, and their pharmaceutically acceptable salts.

Especially relevant to this aspect of the invention is the inhibition of the replication of viruses selected from the group consisting of picorna-, toga-, bunya, orthomyxo-, paramyxo-, rhabdo-, retro-, arena-, hepatitis B-, hepatitis 10 C-, hepatitis D-, adeno-, vaccinia-, papilloma-, herpes-, corona-, varicella- and zoster-virus, in particular human immunodeficiency virus (HIV). Synergistic activity of the pharmaceutical compositions or combined preparations of this invention against viral infection may be readily determined by means of one or more tests such as, but not limited to, the isobologram method, as previously 15 described by Elion et al. in *J. Biol. Chem.* (1954) 208:477-488 and by Baba et al. in *Antimicrob. Agents Chemother.* (1984) 25:515-517, using EC_{50} for calculating the fractional inhibitory concentration (hereinafter referred as FIC). When the minimum FIC index corresponding to the FIC of combined compounds (e.g., $FIC_x + FIC_y$) is equal to 1.0, the combination is said to be 20 additive; when it is between 1.0 and 0.5, the combination is defined as subsynergistic, and when it is lower than 0.5, the combination is by defined as synergistic. When the minimum FIC index is between 1.0 and 2.0, the combination is defined as subantagonistic and, when it is higher than 2.0, the combination is defined as antagonistic.

25 The pharmaceutical composition or combined preparation with synergistic activity against viral infection according to this invention may contain the pteridine compound of formula (II) over a broad content range depending on the contemplated use and the expected effect of the preparation. Generally, the pteridine content of the combined preparation is within the range of 0.1 to 30 99.9% by weight, preferably from 1 to 99% by weight, more preferably from 5 to 95% by weight.

The pharmaceutical compositions and combined preparations according to this invention may be administered orally or in any other suitable

fashion. Oral administration is preferred and the preparation may have the form of a tablet, aqueous dispersion, dispersable powder or granule, emulsion, hard or soft capsule, syrup, elixir or gel. The dosing forms may be prepared using any method known in the art for manufacturing these pharmaceutical compositions and may comprise as additives sweeteners, flavoring agents, coloring agents, preservatives and the like. Carrier materials and excipients are detailed hereinbelow and may include, *inter alia*, calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, binding agents and the like.

10 The pharmaceutical composition or combined preparation of this invention may be included in a gelatin capsule mixed with any inert solid diluent or carrier material, or has the form of a soft gelatin capsule, in which the ingredient is mixed with a water or oil medium. Aqueous dispersions may comprise the biologically active composition or combined preparation in

15 combination with a suspending agent, dispersing agent or wetting agent. Oil dispersions may comprise suspending agents such as a vegetable oil. Rectal administration is also applicable, for instance in the form of suppositories or gels. Injection (e.g. intramuscularly or intraperitoneously) is also applicable as a mode of administration, for instance in the form of injectable solutions or

20 dispersions, depending upon the disorder to be treated and the condition of the patient.

Auto-immune disorders to be prevented or treated by the pharmaceutical compositions or combined preparations of this invention include both systemic auto-immune diseases such as but not limited to lupus

25 erythematosus, psoriasis, vasculitis, polymyositis, scleroderma, multiple sclerosis, ankylosing spondylitis, rheumatoid arthritis and Sjögren syndrome; auto-immune endocrine disorders such as thyroiditis; and organ-specific auto-immune diseases such as but not limited to Addison disease, hemolytic or pernicious anemia, Goodpasture syndrome, Graves disease, idiopathic

30 thrombocytopenic purpura, insulin-dependent diabetes mellitus, juvenile diabetes, uveitis, Crohn's disease, ulcerative colitis, pemphigus, atopic dermatitis, autoimmune hepatitis, primary biliary cirrhosis, autoimmune

pneumonitis, autoimmune carditis, myasthenia gravis, glomerulonephritis and spontaneous infertility.

Transplant rejections to be prevented or treated by the pharmaceutical compositions or combined preparations of this invention include the rejection
5 of transplanted or grafted organs or cells (both allografts and xenografts), such as but not limited to host versus graft reaction disease. The term "organ" as used herein means all organs or parts of organs in mammals, in particular humans, such as but not limited to kidney, lung, bone marrow, hair, cornea, eye (vitreous), heart, heart valve, liver, pancreas, blood vessel, skin, muscle,
10 bone, intestine or stomach. "Rejection" as used herein mean all reactions of the recipient body or of the transplanted organ which in the end lead to cell or tissue death in the transplanted organ or adversely affect the functional ability and viability of the transplanted organ or the recipient. In particular, this means acute and chronic rejection reactions. Also included in this invention is
15 preventing or treating the rejection of cell transplants and xenotransplantation. The major hurdle for xenotransplantation is that even before the T lymphocytes, responsible for the rejection of allografts, are activated, the innate immune system, especially T-independent B lymphocytes and macrophages are activated. This provokes two types of severe and early
20 acute rejection called hyper-acute rejection and vascular rejection, respectively. The present invention addresses the problem that conventional immunosuppressant drugs like cyclosporin A are ineffective in xenotransplantation. The ability of the compounds of this invention to suppress T-independent xeno-antibody production as well as macrophage
25 activation may be evaluated in the ability to prevent xenograft rejection in athymic, T-deficient mice receiving xenogenic hamster-heart grafts.

Cell proliferative disorders to be prevented or treated by the pharmaceutical compositions or combined preparations of this invention include any kind of tumor progression or invasion or metastasis inhibition of a
30 cancer, preferably one selected from the group consisting of lung cancer, leukaemia, ovarian cancer, sarcoma, Kaposi's sarcoma, meningioma, colon cancer, lymph node tumor, glioblastoma multiforme, prostate cancer or skin carcinose.

CNS disorders to be prevented or treated by the pharmaceutical compositions of this invention include cognitive pathologies such as dementia, cerebral ischemia, trauma, epilepsy, schizophrenia, chronic pain and neurologic disorders such as but not limited to depression, social phobia and
5 obsessive compulsive disorders.

Cardiovascular disorders to be prevented or treated by the pharmaceutical compositions of this invention include ischemic disorders, infarct or reperfusion damage, atherosclerosis and stroke.

Allergic conditions to be prevented or treated by the pharmaceutical
10 compositions of this invention include those caused by the pollen of gramineae, the presence of pets, as well as more severe forms, such as asthma, characterized by inflammation of airways and bronchospasm. Without wishing to be bound by theory, the antiallergic effect of the compounds of the invention may be related to their suppression of certain B-cell activation pathways,
15 which can lead to the suppression of IgE release. It may also be related to their properties of inhibiting certain Th2 cytokines, such as IL-5, IL-13 or IL-10, involved in asthma.

The term " pharmaceutically acceptable carrier or excipient " as used herein in relation to pharmaceutical compositions and combined preparations
20 means any material or substance with which the active principle, i.e. the pteridine derivative of formula (II), and optionally the immunosuppressant or immunomodulator or antineoplastic drug or antiviral agent, may be formulated in order to facilitate its application or dissemination to the locus to be treated, for instance by dissolving, dispersing or diffusing the said composition, and/or
25 to facilitate its storage, transport or handling without impairing its effectiveness. The pharmaceutically acceptable carrier may be a solid or a liquid or a gas which has been compressed to form a liquid, i.e. the compositions of this invention can suitably be used as concentrates, emulsions, solutions, granulates, dusts, sprays, aerosols, pellets or powders.

30 Suitable pharmaceutical carriers for use in the said pharmaceutical compositions and their formulation are well known to those skilled in the art. There is no particular restriction to their selection within the present invention although, due to the usually low or very low water-solubility of the pteridine

derivatives of this invention, special attention will be paid to the selection of suitable carrier combinations that can assist in properly formulating them in view of the expected time release profile. Suitable pharmaceutical carriers include additives such as wetting agents, dispersing agents, stickers, adhesives, emulsifying or surface-active agents, thickening agents, complexing agents, gelling agents, solvents, coatings, antibacterial and antifungal agents (for example phenol, sorbic acid, chlorobutanol), isotonic agents (such as sugars or sodium chloride) and the like, provided the same are consistent with pharmaceutical practice, i.e. carriers and additives which do not create permanent damage to mammals. The pharmaceutical compositions of the present invention may be prepared in any known manner, for instance by homogeneously mixing, dissolving, spray-drying, coating and/or grinding the active ingredients, in a one-step or a multi-steps procedure, with the selected carrier material and, where appropriate, the other additives such as surface-active agents. may also be prepared by micronisation, for instance in view to obtain them in the form of microspheres usually having a diameter of about 1 to 10 μm , namely for the manufacture of microcapsules for controlled or sustained release of the biologically active ingredient(s).

Suitable surface-active agents to be used in the pharmaceutical compositions of the present invention are non-ionic, cationic and/or anionic materials having good emulsifying, dispersing and/or wetting properties. Suitable anionic surfactants include both water-soluble soaps and water-soluble synthetic surface-active agents. Suitable soaps are alkaline or alkaline-earth metal salts, unsubstituted or substituted ammonium salts of higher fatty acids ($\text{C}_{10}\text{-C}_{22}$), e.g. the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures obtainable from coconut oil or tallow oil. Synthetic surfactants include sodium or calcium salts of polyacrylic acids; fatty sulphonates and sulphates; sulphonated benzimidazole derivatives and alkylarylsulphonates. Fatty sulphonates or sulphates are usually in the form of alkaline or alkaline-earth metal salts, unsubstituted ammonium salts or ammonium salts substituted with an alkyl or acyl radical having from 8 to 22 carbon atoms, e.g. the sodium or calcium salt of lignosulphonic acid or

dodecylsulphonic acid or a mixture of fatty alcohol sulphates obtained from natural fatty acids, alkaline or alkaline-earth metal salts of sulphuric or sulphonic acid esters (such as sodium lauryl sulphate) and sulphonic acids of fatty alcohol/ethylene oxide adducts. Suitable sulphonated benzimidazole derivatives preferably contain 8 to 22 carbon atoms. Examples of alkylarylsulphonates are the sodium, calcium or alcanolamine salts of dodecylbenzene sulphonic acid or dibutyl-naphtalenesulphonic acid or a naphtalene-sulphonic acid/formaldehyde condensation product. Also suitable are the corresponding phosphates, e.g. salts of phosphoric acid ester and an adduct of p-nonylphenol with ethylene and/or propylene oxide, or phospholipids. Suitable phospholipids for this purpose are the natural (originating from animal or plant cells) or synthetic phospholipids of the cephalin or lecithin type such as e.g. phosphatidylethanolamine, phosphatidylserine, phosphatidylglycerine, lysolecithin, cardiolipin, dioctanyl-phosphatidylcholine, dipalmitoylphosphatidylcholine and their mixtures.

Suitable non-ionic surfactants include polyethoxylated and polypropoxylated derivatives of alkylphenols, fatty alcohols, fatty acids, aliphatic amines or amides containing at least 12 carbon atoms in the molecule, alkylarenesulphonates and dialkylsulphosuccinates, such as polyglycol ether derivatives of aliphatic and cycloaliphatic alcohols, saturated and unsaturated fatty acids and alkylphenols, said derivatives preferably containing 3 to 10 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenol. Further suitable non-ionic surfactants are water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethyleneglycol ether groups and/or 10 to 100 propyleneglycol ether groups. Such compounds usually contain from 1 to 5 ethyleneglycol units per propyleneglycol unit. Representative examples of non-ionic surfactants are nonylphenol-polyethoxyethanol, castor oil polyglycolic ethers, polypropylene/ polyethylene oxide adducts, tributylphenoxypolyethoxyethanol, polyethyleneglycol and octylphenoxypolyethoxyethanol. Fatty acid esters of polyethylene sorbitan

(such as polyoxyethylene sorbitan trioleate), glycerol, sorbitan, sucrose and pentaerythritol are also suitable non-ionic surfactants.

Suitable cationic surfactants include quaternary ammonium salts, preferably halides, having 4 hydrocarbon radicals optionally substituted with
5 halo, phenyl, substituted phenyl or hydroxy; for instance quaternary ammonium salts containing as N-substituent at least one C₈-C₂₂ alkyl radical (e.g. cetyl, lauryl, palmityl, myristyl, oleyl and the like) and, as further substituents, unsubstituted or halogenated lower alkyl, benzyl and/or hydroxy-lower alkyl radicals.

10 A more detailed description of surface-active agents suitable for this purpose may be found for instance in "McCutcheon's Detergents and Emulsifiers Annual" (MC Publishing Corp., Ridgewood, New Jersey, 1981), "Tensid-Taschenbuch", 2nd ed. (Hanser Verlag, Vienna, 1981) and "Encyclopaedia of Surfactants (Chemical Publishing Co., New York, 1981).

15 Structure-forming, thickening or gel-forming agents may be included into the pharmaceutical compositions and combined preparations of the invention. Suitable such agents are in particular highly dispersed silicic acid, such as the product commercially available under the trade name Aerosil; bentonites; tetraalkyl ammonium salts of montmorillonites (e.g., products commercially
20 available under the trade name Bentone), wherein each of the alkyl groups may contain from 1 to 20 carbon atoms; cetostearyl alcohol and modified castor oil products (e.g. the product commercially available under the trade name Antisettle).

Gelling agents which may be included into the pharmaceutical
25 compositions and combined preparations of the present invention include, but are not limited to, cellulose derivatives such as carboxymethylcellulose, cellulose acetate and the like; natural gums such as arabic gum, xanthum gum, tragacanth gum, guar gum and the like; gelatin; silicon dioxide; synthetic polymers such as carbomers, and mixtures thereof. Gelatin and modified
30 celluloses represent a preferred class of gelling agents.

Other optional excipients which may be included in the pharmaceutical compositions and combined preparations of the present invention include

additives such as magnesium oxide; azo dyes; organic and inorganic pigments such as titanium dioxide; UV-absorbers; stabilisers; odor masking agents; viscosity enhancers; antioxidants such as, for example, ascorbyl palmitate, sodium bisulfite, sodium metabisulfite and the like, and mixtures thereof; preservatives such as, for example, potassium sorbate, sodium benzoate, sorbic acid, propyl gallate, benzylalcohol, methyl paraben, propyl paraben and the like; sequestering agents such as ethylene-diamine tetraacetic acid; flavoring agents such as natural vanillin; buffers such as citric acid and acetic acid; extenders or bulking agents such as silicates, diatomaceous earth, magnesium oxide or aluminum oxide; densification agents such as magnesium salts; and mixtures thereof.

Additional ingredients may be included in order to control the duration of action of the biologically-active ingredient in the compositions and combined preparations of the invention. Control release compositions may thus be achieved by selecting appropriate polymer carriers such as for example polyesters, polyamino-acids, polyvinyl-pyrrolidone, ethylene-vinyl acetate copolymers, methylcellulose, carboxymethylcellulose, protamine sulfate and the like. The rate of drug release and duration of action may also be controlled by incorporating the active ingredient into particles, e.g. microcapsules, of a polymeric substance such as hydrogels, polylactic acid, hydroxymethyl-cellulose, polymethyl methacrylate and the other above-described polymers. Such methods include colloid drug delivery systems like liposomes, microspheres, microemulsions, nanoparticles, nanocapsules and so on. Depending on the route of administration, the pharmaceutical composition or combined preparation of the invention may also require protective coatings.

Pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation thereof. Typical carriers for this purpose therefore include biocompatible aqueous buffers, ethanol, glycerol, propylene glycol, polyethylene glycol, complexing agents such as cyclodextrins and the like, and mixtures thereof.

Since, in the case of combined preparations including the pteridine derivative of this invention and an immunosuppressant or immunomodulator

or antihistamine or antineoplastic drug or antiviral agent, both ingredients do not necessarily bring out their synergistic therapeutic effect directly at the same time in the patient to be treated, the said combined preparation may be in the form of a medical kit or package containing the two ingredients in
 5 separate but adjacent form. In the latter context, each ingredient may therefore be formulated in a way suitable for an administration route different from that of the other ingredient, e.g. one of them may be in the form of an oral or parenteral formulation whereas the other is in the form of an ampoule for intravenous injection or an aerosol.

10 The present invention further relates to a method for preventing or treating a disease selected from the group consisting of CNS disorders, cell proliferative disorders, allergic conditions, viral infections, immune and auto-immune disorders and transplant rejections in a subject or patient by administering to the patient in need thereof an effective amount of a pteridine
 15 derivative having the general formula (II), optionally together with an effective amount of another immunosuppressant or immunomodulator or antineoplastic drug or antiviral agent, or a pharmaceutical composition such as disclosed above in extensive details. The effective amount is usually in the range of 0.01 mg to 20 mg, preferably 0.1 mg to 5 mg, per day per kg bodyweight for
 20 humans. Depending upon the pathologic condition to be treated and the patient's condition, the said effective amount may be divided into several sub-units per day or may be administered at more than one day intervals. The patient to be treated may be any warm-blooded animal, preferably a human being, suffering from said pathologic condition.

25 The following examples are intended to illustrate and not to limit the scope of the present invention in all its aspects.

Example 1 – preparation of 2-amino-4-n-pentvloxv-6-styrylpteridine.

30 A mixture of 1.5 g (5.6 mmoles) 2-amino-6-chloro-4-n-pentyloxypteridine (e.g. available following the procedure disclosed by Mohr et al. in *Helv. Chem. Acta* (1992) 75:2317), palladium acetate (63 mg, 0.28 mmoles), tri-o-tolylphosphane (682 mg, 2.24 mmoles), cuprous iodide (53 mg, 0.28 mmoles), styrene (1,3 ml., 11.3 mmoles) and triethylamine 3.1 ml, 22 mmoles) was

stirred in dry acetonitrile (50 ml) under reflux for 90 hours. It was evaporated and the residue purified by silica gel column chromatography with chloroform. The product fraction was evaporated to give 1.37 g (yield: 72%) of an orange powder exhibiting, after recrystallization from a EtOAc/hexane mixture, a melting point (m.p.) range of 127-128 °C.

Example 2 – preparation of 2-amino-6-(1,2-dibromophenethyl)-4-n-pentyloxy-pteridine.

To a solution of the derivative of example 1 (1.0 g, 2.94 mmoles) in chloroform (50 ml) was added a 2 M bromine solution in chloroform (2.2 ml, 4.4 mmoles) and then the mixture was stirred at room temperature for 7 hours. It was diluted with chloroform (50 ml), washed with a saturated aqueous Na₂SO₃ solution (100 ml) and dried over sodium sulfate. After evaporation of the solvents, the residue was treated with toluene, filtered, washed with ether and dried in a vacuum desiccator to give 0.84 g (yield: 57%) of a yellow powder.

Example 3 – preparation of 2-amino-4,7-dimethoxy-6-styrylpteridine.

A suspension of the derivative of example 2 (0.3 g, 0.6 mmoles) in methanol (10 ml) was treated with 1 M methanolic sodium methoxide (3 ml, 3 mmoles) and then refluxed for 4 hours. It was diluted with chloroform (100 ml), washed with a saturated aqueous ammonium chloride solution and water and then the solution was dried over sodium sulfate. The filtrate was evaporated and the residue was purified by silica gel column chromatography while using chloroform as the eluent. The product fraction was evaporated to give 50 mg (yield: 26%) of a yellow powder with a melting point range of 197-198 °C.

Example 4 – preparation of 0⁴-methyl-biopterin (2-amino-4-methoxy-6-(1,2-dihydroxypropyl) pteridine)

To a solution of N², 1', 2'-O-triacetyl-biopterin (1,0 g, 2.75 mmoles), triphenyl-phosphane (12,08 g, 4.13 mmoles) and methanol (0.15 ml, 3.7 mmoles) in dry dioxane (30 ml) was added diisopropyl azodicarboxylate (0.81 g, 4.1 mmoles). After stirring for 1.5 hour at room temperature, the mixture

was evaporated to dryness. The residue was purified by silica gel column chromatography while using an ethylacetate/ CHCl_3 (1:4) mixture as the eluent. The product fraction was evaporated and dried in vacuum to give 0.4 g (yield: 38%) of $\text{N}^2,1',2'$ -O-triacetyl- O^4 -methylbiopterin. Deacetylation of this reaction product (0.28 g, 0.74 mmole) was done by stirring it in absolute methanol (20 ml) and triethylamine (4 ml) for 24 hours. Evaporation to dryness, treatment of the residue with ether, filtration and drying gave 0.172 g (yield: 83%) of O^4 -methyl-biopterin with a melting point range of 160-161 °C.

10 Example 5 – preparation of 2-amino-4-hydroxylamino-6-phenylpteridine.

A suspension of 2,5,6-triamino-4-methoxypyrimidine dihydrochloride (1 g, 4 mmoles) in methanol (40 ml) was heated to boiling and then a solution of phenylglyoxalmonoxime (1 g, 6.6 mmoles) in methanol (10 ml) was added dropwise. A clear solution is obtained from which on reflux for 2 hours a precipitate was separated out. The solid (hydrochloride salt) was filtered off, suspended in water (30 ml) and then neutralized to pH 8 by concentrated ammonia. The resulting precipitate was collected, washed with water and ethanol and dried at 100 °C to give 0.84 g of a yellow powder (yield: 82%).

20 Examples 6 to 53 - synthesis of 2-amino-4-dialkylamino-6-arylpteridines, 2-amino-4-di(arylalkyl)amino-6-arylpteridines, 2-amino-4-alkylamino-6-arylpteridines, 2-amino-4-(N-containing heterocyclic amino)-6-arylpteridines and 2-amino-4-alkoxy-6-arylpteridines

The procedure for the synthesis of the following 2-amino-4-dialkylamino-6-arylpteridines, 2-amino-4-dialkylamino-6-arylpteridines, 2-amino-4-alkylamino-6-arylpteridines, 2-amino-4-(N-containing heterocyclic amino)-6-arylpteridines and 2-amino-4-alkoxy-6-arylpteridines proceeds in three steps:

a) a solution of 2,6-diamino-4-chloro-5-p-chlorophenylazopyrimidine (a compound known from British Patent No. 677,342) (5.0 g, 16.6 mmoles) in DMF (50ml) and 0.12 mole of the appropriate reactant, being selected from the group consisting of secondary alkylamines and arylalkylamines (e.g. dimethyl-amine in ethanol (50%), diethylamine, di-n-propylamine or dibenzylamine), primary amines (e.g. an adamantanamine), heterocyclic

- amines (e.g. morpholine, piperidine, pyrrolidine, piperazine or N-methyl-piperazine) and alkaline metal alkoxides (e.g. sodium ethoxide or sodium isopropoxide), were heated in an oil bath at 70°C for 5 hours. Then water (50 ml) was added, cooled and the yellow precipitate collected, washed with water and dried. Recrystallization from ethanol or a DMF/water mixture provided the relevant 2,6-diamino-4-dialkylamino-5-p-chlorophenylazopyrimidine, 2,6-diamino-4-di(arylalkyl)amino-5-p-chlorophenylazopyrimidine, 2,6-diamino-4-alkyl-amino-5-p-chlorophenylazopyrimidine, 2,6-diamino-4-(N-containing hetero-cyclic amino)-5-p-chlorophenylazopyrimidine or 2,6-diamino-4-alkoxy-5-p-chlorophenylazopyrimidine with a yield ranging from 55 to 90%.
- b) a suspension of the pyrimidine compound (3.28 g, 10 mmoles) resulting from step (a) in methanol (70 ml) and concentrated ammonia (10 ml) was reduced in a shaking apparatus under a hydrogen atmosphere in the presence of a Raney nickel catalyst (3.5 g) for 2 days. The catalyst was filtered off under argon atmosphere and then the filtrate evaporated in vacuo to dryness. The residue was treated with ether to remove p-chloroaniline, filtered and then the solid stirred in methanolic HCl (10%, 50 ml) overnight. The dihydrochloride salt (obtained with a yield ranging from 85 to 90%) of the relevant 2,5,6-triamino-4-dialkylaminopyrimidine, 2,5,6-triamino-4-alkoxypyrimidine, 2,5,6-triamino-4-di(arylalkyl)aminopyrimidine, 2,5,6-triamino-4-alkylaminopyrimidine or 2,5,6-triamino-4-(N-containing heterocyclic amino) pyrimidine, was collected and dried in a vacuum desiccator over KOH.
- c) to a boiling solution of the 2,5,6-triamino-4-substituted pyrimidine dihydrochloride salt (5 mmoles) from step (b) in methanol (20 ml) was added a solution of the relevant arylglyoxalmonoxime (7.5 mmoles) in methanol (10 ml) dropwise and then the mixture was heated under reflux for 3 hours. After cooling, the suspension or solution was made alkaline by means of concentrated ammonia up to pH 9 and the resulting precipitate was filtered off, washed with water and dried. Recrystallization was done from ethanol and a DMF/water mixture, respectively, such as to provide a yellow solid with a yield ranging from 50 to 85%.

The following compounds were prepared according to the above general procedure:

- 2-amino-4-dimethylamino-6-phenylpteridine (example 6);
- 2-amino-4-dimethylamino-6-(4-tolyl) pteridine (example 7);
- 5 2-amino-4-dimethylamino-6-(4-methoxyphenyl)pteridine (example 8);
- 2-amino-4-diethylamino-6-phenylpteridine (example 9);
- 2-amino-4-diethylamino-6-(4-chlorophenyl)pteridine (example 10);
- 2-amino-4-diethylamino-6-(4-methoxyphenyl)pteridine (example 11);
- 2-amino-4-diethylamino-6-(3,4-dimethoxyphenyl)pteridine (example 12);
- 10 2-amino-4-dibenzylamino-6-phenylpteridine (example 13);
- 2-amino-4-dibenzylamino-6-(4-chlorophenyl)pteridine (example 14);
- 2-amino-4-dibenzylamino-6-(4-methoxyphenyl)pteridine (example 15);
- 2-amino-4-dibenzylamino-6-(3,4-dimethoxyphenyl)pteridine (example 16);
- 2-amino-4-dipropylamino-6-phenylpteridine(example 17);
- 15 2-amino-4-dipropylamino-6-(4-chlorophenyl)pteridine (example 18);
- 2-amino-4-dipropylamino-6-(4-methoxyphenyl)pteridine (example 19);
- 2-amino-4-dipropylamino-6-(3,4-dimethoxyphenyl)pteridine (example 20);
- 2-amino-4-morpholino-6-phenylpteridine (example 21);
- 2-amino-4-morpholino-6-(4-chlorophenyl)pteridine (example 22);
- 20 2-amino-4-morpholino-6-(4-methoxyphenyl)pteridine (example 23);
- 2-amino-4-morpholino-6-(3,4-dimethoxyphenyl)pteridine (example 24);
- 2-amino-4-piperidino-6-phenylpteridine (example 25);
- 2-amino-4-piperidino-6-(4-chlorophenyl)pteridine (example 26);
- 2-amino-4-piperidino-6-(4-methoxyphenyl)pteridine (example 27);
- 25 2-amino-4-piperidino-6-(3,4-dimethoxyphenyl)pteridine (example 28);
- 2-amino-4-N-methylpiperazino-6-phenylpteridine (example 29);
- 2-amino-4-N-methylpiperazino-6-(4-chlorophenyl)pteridine (example 30);
- 2-amino-4-N-methylpiperazino-6- (4-methoxyphenyl) pteridine (example 31);
- 2-amino-4-methylpiperazino-6- (3, 4-dimethoxyphenyl) pteridine (example 32);
- 30 2-amino-4-pyrrolidino-6- (4-methoxyphenyl) pteridine (example 33);
- 2-amino-4-piperazino-6-phenylpteridine (example 34);
- 2-amino-4-piperazino-6- (4-chlorophenyl) pteridine (example 35);
- 2-amino-4-piperazino-6- (4-methoxyphenyl) pteridine (example 36);

- 2-amino-4-piperazino-6- (3, 4-dimethoxyphenyl) pteridine (example 37);
 2-amino-4-dibenzylamino-6-(3, 4,5-trimethoxyphenyl) pteridine (example 38);
 2-amino-4-morpholino-6- (3, 4, 5-trimethoxyphenyl) pteridine (example 39);
 2-amino-4-(3-adamantylamino)-6-(3,4,5-trimethoxyphenyl) pteridine (example
 5 40);
 2-amino-4-(3-adamantylamino)-6-naphtylpteridine (example 41);
 2-amino-4-(4-adamantylamino)-6-(3,4,5-trimethoxyphenyl) pteridine (example
 42);
 2-amino-4-(4-adamantylamino)-6-naphtylpteridine (example 43);
 10 2-amino-4-morpholino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine
 (example 44);
 2-amino-4-dimethylamino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine
 (example 45);
 2-amino-4-pyrrolidino-6-(3, 4-dimethoxyphenyl) pteridine (example 46);
 15 2-amino-4-dimethylamino-6-(3, 4-dimethoxyphenyl) pteridine (example 47);
 2-amino-4-dimethylamino-6-methylpteridine (example 48);
 2-amino-4-ethoxy-6-phenylpteridine (example 49);
 2-amino-4-propylamino-6-phenylpteridine (example 50);
 2-amino-4-propylamino-6-(3, 4-dimethoxyphenyl) pteridine (example 51);
 20 2-acetamido-4-isopropoxy-6-(3,4-dimethoxyphenyl) pteridine (example 52);
 and
 2-amino-4-ethoxy-6-(3,4-dimethoxyphenyl)pteridine (example 53);

Example 54 - synthesis of 2,6-diamino-4-ethoxy-pyrimidine

- 25 To a solution of sodium (1.05 g) in ethanol (50 ml) was added 4-chloro-
 2,6-diaminopyrimidine (6 g, 41.4 mmoles). The resulting solution was heated
 in a reactor for 6 hours at 160 °C. The reaction mixture was cooled down and
 the precipitated sodium chloride was filtered off. The filtrate was concentrated
 and precipitated from ethanol (two times), affording the pure title compound as
 30 a white solid (4.53 g, 72% yield). The spectral data are identical to those
 described e.g. by W. Pfeiderer et al. in *Chem. Ber.* (1961) 94, 12.

Example 55 - synthesis of 2,6-diamino-4-isopropoxy-pyrimidine

The same procedure as in example 54 was followed using isopropanol instead of ethanol. The filtrate was pure enough for further reaction without purification. The spectral data are identical to those described e.g. by W.

5 Pfeiderer et al. in *Chem. Ber.* (1961) 94, 12.

Example 56 - synthesis of 5-nitroso-2,6-diamino-4-ethoxy-pyrimidine

To a solution of the compound of example 54 (6.13 g, 39.8 mmoles) in 20 % aqueous acetic acid (57 ml) was added dropwise a solution of NaNO₂ (3.29 g) in water (13 ml) at 80 °C. A pink precipitate was formed which was stirred at 80°C for an additional 2 hours. The reaction mixture was cooled down in the refrigerator overnight and the resulting precipitate was filtered off, yielding the title compound as a pink powder (4.98 g, yield 68 %). Spectral data are identical with those described e.g. by W. Pfeiderer et al. in *Chem.*

10 *Ber.* (1961) 94, 12.

Example 57 - synthesis of 5-nitroso-2,6-diamino-4-isopropoxy-pyrimidine

The same procedure was followed as in example 56 but starting from the compound of example 55. The product has identical spectral data to those described by W. Pfeiderer et al. (cited *supra*).

20

Example 58 - synthesis of 2,5,6-triamino-4-ethoxy-pyrimidine

To a suspension of the compound of example 56 (7.12 g, 38.9 mmoles) in water (150 ml) at 60 °C was added sodium dithionite (46.7 mmol, 8.12 g). Additional sodium dithionite was added till the pink colour completely disappeared and a yellow solution was formed. The solution was stirred at 60 °C for another 4 hours. Water was evaporated and the resulting residue was precipitated from a small amount of water, providing the title compound as a yellow powder (4.02 g, yield 61 %). Spectral data are identical with literature data (W. Pfeiderer et al. cited *supra*).

25
30

Example 59 - synthesis of 2,5,6-triamino-4-isopropoxy-pyrimidine

The procedure of example 58 was followed, however using the compound of example 57 as the starting material. The spectral data of the product obtained are identical with the literature data (W. Pfeiderer et al. cited *supra*).

Example 60 - synthesis of 2-amino-4-ethoxy-pteridin

To a solution of 2,5,6-triamino-4-ethoxy-pyrimidine (10.54 g, 62.37 mmoles) in ethanol (160 ml) was added glyoxal (40 % solution in water, 2.7 ml, 18.6 mmoles). The reaction mixture was refluxed for 4 hours. Some insoluble material was filtered off. The filtrate was concentrated *in vacuo* and the residue purified by flash chromatography (silica, using a CH₃OH/CH₂Cl₂ mixture (5:95) as the eluent), providing the pure title compound (7.34 g, yield: 62 %). The spectral data of the product are identical with the literature data (W. Pfeiderer et al. cited *supra*).

Example 61 - synthesis of 2-amino-4-isopropoxy-pteridin

The procedure of example 60 was repeated, however using isopropanol as the solvent instead of ethanol. The spectral data of the product obtained are identical with the literature data (W. Pfeiderer et al. cited *supra*).

Example 62 - synthesis of 2-amino-4-ethoxypteridine-N⁸-oxide

To a cooled (0 °C) solution of the compound of example 60 (2.47 g, 12.9 mmoles) in trifluoroacetic acid (53 ml) was added dropwise 2.53 ml of a 35 % aqueous H₂O₂ solution. The reaction mixture was kept at 4 °C for two days in the refrigerator, whereby another 1.25 ml of the same H₂O₂ solution was added after 1 day. The solution was concentrated *in vacuo*. The residue was suspended in water and neutralized by the addition of a concentrated ammonia solution. Evaporation of the solvent *in vacuo* and purification of the residue by flash chromatography (silica, using a CH₃OH/CH₂Cl₂ mixture (6:94) as the eluent) provided the title compound as a yellow powder (861 mg, yield: 32 %). Mass spectrum data are as follows: *m/z* (%): 230 ([M+Na]⁺, 30), 208 ([M+H]⁺, 100), 180 [(M+H-ethene)⁺, 10].

Example 63 - synthesis of 2-amino-4-isopropoxypteridine-N⁸-oxide

The procedure as described in example 62 was followed, however using the compound of example 61 as the starting material. Mass spectrum data are as follows: *m/z* (%): 222 ([M+H]⁺, 100), 180 ([M+H-propene]⁺, 60)

5

Example 64 - synthesis of 2-amino-6-chloro-4-ethoxypteridine

A suspension of the compound of example 62 (460 mg, 2.22 mmoles) in acetyl chloride (5.5 ml) was stirred at -40 °C. Trifluoroacetic acid (1.69 ml) was then added dropwise. The resulting solution was slowly warmed up to 0 °C and stirred for an additional 4 hours at 0 °C. Reaction was carefully quenched with ice, followed by neutralization with a concentrated ammonia solution (pH = 8). The aqueous phase was extracted with CH₂Cl₂ (five times). The combined organic layers were concentrated *in vacuo* and the residue was purified by flash chromatography (silica, using a CH₃OH/CH₂Cl₂ mixture (1:99) as the eluent), thus providing the title compound as a yellow powder (360 mg, yield: 72 %). This compound was further characterized as follows:

- mass spectrum: *m/z* (%): 226 ([M+H]⁺, 100),
- ¹H-NMR (200 MHz, DMSO-*d*₆): δ 1.42 (3 H, t), 4.52 (2 H, q), 7.42 (2 H, d) and 8.85 (1 H, s) ppm,
- ¹³C-NMR (50 MHz, DMSO-*d*₆): δ 14.19, 63.58, 121.74, 140.22, 150.99, 156.13, 161.98 and 165.97 ppm.

20

Example 65 - synthesis of 2-amino-6-chloro-4-isopropoxypteridine

The procedure as described in example 64 was followed, however starting from the compound of example 63. The mass spectrum data of the resulting compound are as follows: *m/z* (%): 240 ([M+H]⁺, 55), 198 ([M+H-propene]⁺, 100).

25

Examples 66 to 83 - synthesis of 2-amino-6-aryl-4-ethoxypteridines and 2-amino-6-heteroaryl-4-ethoxypteridines

30

The general procedure used for preparing 2-amino-6-aryl-4-ethoxypteridines is as follows: to a degassed solution of the compound of example 64 (50 mg, 0.22 mmole) in THF (5 ml) was added a degassed solution of

sodium carbonate (5 ml of a 0.4 M solution in water), tetrakis(triphenylphosphine) palladium (0.013 mmole, 14 mg) and an arylboronic or (examples 72 and 73) heteroarylboronic acid (0.22 mmole). The solution was refluxed for 4 hours. Solvents were concentrated *in vacuo* and the residue was purified by
 5 flash chromatography (silica) with an appropriate CH₃OH/CH₂Cl₂ mixture (2:98 or 3:97) as the eluent (except for the compound of example 82, which was eluted with an acetone/CH₂Cl₂ (7:3) mixture). This procedure provided, with a yield ranging from 16 % to 60 % depending upon the aryl or heteroaryl group (from the arylboronic or heteroarylboronic acid) introduced at the 6-position of
 10 the pteridine ring, the following pure final compounds which were characterized by their mass spectrum MS and optionally by their ¹H-NMR (200 MHz, DMSO-*d*₆) spectrum:

- 2-amino-6-(p-methoxyphenyl)-4-ethoxy-pteridine (example 66): MS 298 ([M+H]⁺, 100), 270 ([M+H-ethene]⁺, 55);
- 15 - 2-amino-6-(o-methoxyphenyl)-4-ethoxy-pteridine (example 67): MS 298 ([M+H]⁺, 100), 270 ([M+H-ethene]⁺, 30);
- 2-amino-6-(m-methoxyphenyl)-4-ethoxy-pteridine (example 68): MS 298 ([M+H]⁺, 100), 270 ([M+H-ethene]⁺, 35); ¹H-NMR: 1.46 (3 H, t), 3.85 (3 H, s), 4.58 (2 H, q), 7.06 (1 H, dd), 7.33 (2 H, br s), 7.46 (1 H, t),
 20 7.68 (1 H, m) and 9.43 (1 H, s) ppm;
- 2-amino-6-(3,4-difluorophenyl)-4-ethoxy-pteridine (example 69): MS 304 ([M+H]⁺, 100), 270 ([M+H-ethene]⁺, 35); ¹H-NMR: 1.45 (3 H, t), 4.57 (2 H, q), 7.42 (2 H, br s), 7.60 (1 H, q), 7.98 (1 H, d), 8.16 (1 H, t) and 9.42 (1 H, s) ppm;
- 25 - 2-amino-6-(p-dimethylaminophenyl)-4-ethoxy-pteridine (example 70): MS 311 ([M+H]⁺, 100), 283 ([M+H-ethene]⁺, 35);
- 2-amino-6-(p-trifluoromethylphenyl)-4-ethoxy-pteridine (example 71): MS 336 ([M+H]⁺, 100), 308 ([M+H-ethene]⁺, 50);
- 2-amino-6-(2-thienyl)-4-ethoxy-pteridine (example 72): MS 274 ([M+H]⁺,
 30 100), 246 ([M+H-ethene]⁺, 40);
- 2-amino-6-(3-thienyl)-4-ethoxy-pteridine (example 73): MS 274 ([M+H]⁺, 100), 246 ([M+H-ethene]⁺, 45);

- 2-amino-6-(3,4-dichlorophenyl)-4-ethoxy-pteridine (example 74): MS 337 ($[M+H]^+$, 100); 1H -NMR: 1.46 (3 H, t), 4.59 (2 H, q), 7.42 (2 H, br s), 7.81 (1H, d), 8.14 (1 H, dd), 8.37 (1 H, d) and 9.47 (1 H, s) ppm;
- 2-amino-6-(p-cyanophenyl)-4-ethoxy-pteridine (example 75): MS 293 ($[M+H]^+$, 100), 265 ($[M+H-ethene]^+$, 65);
- 2-amino-6-(p-ethoxyphenyl)-4-ethoxy-pteridine (example 76): MS 312 ($[M+H]^+$, 100), 284 ($[M+H-ethene]^+$, 70);
- 2-amino-6-(p-fluorophenyl)-4-ethoxy-pteridine (example 77): MS 286 ($[M+H]^+$, 100), 258 ($[M+H-ethene]^+$, 45);
- 2-amino-6-(p-ethylphenyl)-4-ethoxy-pteridine (example 78): MS 296 ($[M+H]^+$, 100), 268 ($[M+H-ethene]^+$, 45);
- 2-amino-6-(p-acetylphenyl)-4-ethoxy-pteridine (example 79): MS 310 ($[M+H]^+$, 100), 282 ($[M+H-ethene]^+$, 60);
- 2-amino-6-(3-methyl-4-fluorophenyl)-4-ethoxy-pteridine (example 80): MS 300 ($[M+H]^+$, 100), 272 ($[M+H-ethene]^+$, 30);
- 2-amino-6-(p-thiomethylphenyl)-4-ethoxy-pteridine (example 81): MS 314 ($[M+H]^+$, 100), 286 ($[M+H-ethene]^+$, 35);
- 2-amino-6-(p-N,N-dimethylbenzamido)-4-ethoxy-pteridine (example 82) MS 338 ($[M+H]^+$, 100), 311 ($[M+H-ethene]^+$, 15); and
- 2-amino-6-(3,4-dimethoxyphenyl)-4-ethoxy-pteridine (example 83): MS 328 ($[M+H]^+$, 100), 300 ($[M+H-ethene]^+$, 40).

Examples 84 to 98 - synthesis of 2-amino-6-aryl-4-isopropoxypteridines and 2-amino-6-heteroaryl-4-isopropoxypteridines

The procedure as described in examples 66-83 was followed while using 2-amino-6-chloro-4-isopropoxypteridine as the starting material, except that longer reaction times were needed (refluxing overnight instead of 4 hours). This procedure provided, with a yield ranging from 10 % to 70 % depending upon the aryl or heteroaryl group introduced at the 6-position of the pteridine ring, the following pure final compounds which were characterized by their mass spectrum:

- 2-amino-6-(3-methyl-4-methoxyphenyl)-4-isopropoxypteridine (example 84): MS 326 ($[M+H]^+$, 100), 284 ($[M+H-propene]^+$, 30);

- 2-amino-6-(3,4-dimethylphenyl)-4-isopropoxypteridine (example 85): MS 310 ($[M+H]^+$, 100), 268 ($[M+H\text{-propene}]^+$, 60);
- 2-amino-6-(3-chloro-4-trifluoromethylphenyl)-4-isopropoxypteridine (example 86): MS 384 ($[M+H]^+$, 20), 342 ($[M+H\text{-propene}]^+$, 50);
- 5 - 2-amino-6-(3-chloro-4-fluorophenyl)-4-isopropoxypteridine (example 87): MS 334 ($[M+H]^+$, 20), 292 ($[M+H\text{-propene}]^+$, 50);
- 2-amino-6-(p-N,N-diethylbenzamido)-4-isopropoxypteridine (example 88): MS 381 ($[M+H]^+$, 100);
- 2-amino-6-(p-trifluoromethylphenyl)-4-isopropoxypteridine (example 10 89): MS 350 ($[M+H]^+$, 100), 308 ($[M+H\text{-propene}]^+$, 30);
- 2-amino-6-(3,4-difluorophenyl)-4-isopropoxypteridine (example 90): MS 318 ($[M+H]^+$, 100), 276 ($[M+H\text{-propene}]^+$, 50);
- 2-amino-6-(p-methoxyphenyl)-4-isopropoxypteridine (example 91): MS 312 ($[M+H]^+$, 100), 270 ($[M+H\text{-propene}]^+$, 50);
- 15 - 2-amino-6-(p-ethoxyphenyl)-4-isopropoxypteridine (example 92): MS 326 ($[M+H]^+$, 55), 284 ($[M+H\text{-propene}]^+$, 100);
- 2-amino-6-(p-dimethylbenzamido)-4-isopropoxypteridine (example 93): MS 353 ($[M+H]^+$, 75), 311 ($[M+H\text{-propene}]^+$, 100);
- 2-amino-6-(3-thienyl)-4-isopropoxypteridine (example 94): MS 288 20 ($[M+H]^+$, 55), 246 ($[M+H\text{-propene}]^+$, 100);
- 2-amino-6-(p-cyanophenyl)-4-isopropoxypteridine (example 95): MS 307 ($[M+H]^+$, 40), 265 ($[M+H\text{-propene}]^+$, 100);
- 2-amino-6-(p-benzoic acid methyl ester)-4-isopropoxypteridine (example 96): MS 340 ($[M+H]^+$, 75), 298 ($[M+H\text{-propene}]^+$, 100);
- 25 - 2-amino-6-(p-acetylphenyl)-4-isopropoxypteridine (example 97): MS 324 ($[M+H]^+$, 55), 282 ($[M+H\text{-propene}]^+$, 100); and
- 2-amino-6-(3,4-dimethoxyphenyl)-4-isopropoxypteridine (example 98): MS 342 ($[M+H]^+$, 100), 300 ($[M+H\text{-propene}]^+$, 60).

30 Example 99 - synthesis of 2,6-diamino-5-nitroso-4-hydroxypyrimidine

To a solution of 2,6-diamino-4-hydroxypyrimidine (12.9 g, 102.2 mmol) in 200 ml of a 10% acetic acid solution in water at 80 °C was added dropwise a solution of NaNO_2 (7.05 g, 102.2 mmol) in 20 ml water. A pink

precipitate was formed, which was further stirred for 1 hour at 80 °C. The reaction mixture was cooled down in the refrigerator overnight. The precipitate was filtered off and dried over P₂O₅, providing the title compound as a pink powder (15.43 g, yield: 97%). The spectral data are in accordance with literature data (Landauer et al. in *J. Chem. Soc.* (1953) 3721-3722).

Example 100 - synthesis of 2,5,6-triamino-4-hydroxypyrimidine

A suspension of the compound of example 99 (15 g, 96.7 mmoles) in an ammonium sulfide solution (20 % in water, 200 ml) was stirred overnight at 50 °C. The reaction mixture was cooled down in the refrigerator and the precipitate was filtered off, providing the title compound as a yellow powder (11.33 g, yield: 83 %). The spectral data are identical with literature data (Landauer et al. cited *supra*).

Example 101 - synthesis of 2-amino-6-(3,4-dimethoxyphenyl)pterine

To a boiling solution of the compound of example 100 (2.4 g, 17 mmoles) in methanol (100 ml, with 0.9 N HCl) was added dropwise a solution of 3,4-dimethoxyphenylglyoxal mono-oxime (3.8 g, 18 mmoles) in methanol (100 ml). The reaction mixture was heated under reflux for 4 hours. The precipitate formed was filtered off, washed with water, then ethanol and diethyl ether, and dried over P₂O₅ under vacuum, providing the title compound as a yellow powder (4.33 g, yield: 85 %). This compound was further characterized by the following spectra:

- ¹H-NMR (500 MHz, TFA): δ 4.11 (3 H, s), 4.07 (3 H, s), 7.21 (1 H, d), 7.78 (1 H, dd), 7.81 (1 H, d) and 9.32 (1 H, s) ppm;
- ¹³C-NMR (125 MHz, TFA): δ 56.39, 56.7, 111.94, 113.21, 123.22, 127.41, 127.91, 145.92, 149.39, 150.46, 152.47, 153.15, 155.13 and 161.59 ppm.

Example 102 - synthesis of 2-acetylamino-6-(3,4-dimethoxyphenyl)pterine

A suspension of the compound of example 101 (10.46 g, 35 mmoles) in acetic anhydride (600 ml) and acetic acid (200 ml) was refluxed for 1 hour until a clear solution was formed. By cooling down the reaction mixture in the

refrigerator, the precipitate formed was filtered off, washed with ethyl acetate and diethyl ether, and then dried over P_2O_5 under vacuum, providing the title compound as a yellow powder (9.19 g, yield: 77 %). This compound was further characterized by the following spectra:

- 5 - MS: m/z (%): 300 ($[M+H]^+$, 100);
 - 1H -NMR (200 MHz, $DMSO-d_6$): δ 2.22 (3 H, s), 3.84 (3 H, s), 3.87 (3 H, s), 7.14 (1 H, d), 7.75 (2 H, m) and 9.51 (1 H, s) ppm.

10 Example 103 - synthesis of 2-acetylamino-4-(1,2,4-triazolyl)-6-(3,4-dimethoxy-phenyl)pteridine

To a solution of phosphorus oxychloride (1.68 ml, 18 mmoles) and 1,2,4-triazole (4.96 g, 72 mmoles) in dry pyridine (110 ml) was added the compound of example 102 (2.45 g, 7.18 mmoles). The suspension was stirred at room temperature for 4 hours. The precipitate was filtered off, washed with
 15 pyridine, toluene and diethyl ether. The resulting solid was dried over P_2O_5 under vacuum, providing the title compound as a yellow powder (2 g, yield: 80 %) which afforded the following mass spectrum: 392 ($[M+H]^+$, 100).

20 Examples 104 and 105 - synthesis of 2-amino-4-mercaptoethyl-6-(3,4-dimethoxyphenyl)pteridine and 2-amino-4-mercaptoisopropyl-6-(3,4-dimethoxyphenyl) pteridine

To a suspension of the compound of example 103 (0.25 mmole, 100 mg) in dioxane (5 ml) was added 1 mmole of either ethanethiol (example 104) or isopropanethiol (example 105) and sodium (12 mg, 0.5 mmole). The
 25 suspension was stirred for 24 hours at room temperature. The solvent was concentrated *in vacuo* and the residue purified by flash chromatography (silica, using a CH_3OH/CH_2Cl_2 mixture (5:95) as an eluent), followed by purification by preparative TLC, providing the pure title compounds as yellow powders with yields ranging from 20 to 30%. Both compounds were
 30 characterized by their mass spectrum as follows:

- 2-amino-4-mercaptoethyl-6-(3,4-dimethoxyphenyl) pteridine: 344 ($[M+H]^+$, 100);

- 2-amino-4-mercaptoisopropyl-6-(3,4-dimethoxyphenyl) pteridine: 357 ([M+H]⁺, 100).

Example 106 - synthesis of a mixture of 2,4-diamino-6-(p-methoxyphenyl)

5 pteridine and 2,4-diamino-7-(p-methoxyphenyl)pteridine

2,4,5,6-tetra-aminopyrimidine (10 mmoles, 1.4 g) was dissolved in water (50 ml) and the pH was adjusted to 9 with ammonium hydroxide. A solution of 4-methoxyphenylglyoxal (11 mmoles, 1.8 g) in ethanol (10 ml) was added dropwise and the solution was refluxed for 1 hour. The yellow precipitate
10 formed was filtered off and washed with water, ethanol and diethyl ether. NMR analysis reveals the obtention of a mixture (1.2 g, 45 % yield) consisting of 87 % of 2,4-diamino-7-(p-methoxyphenyl)pteridine and 13 % of 2,4-diamino-7-(p-methoxy-phenyl)pteridine. ¹H-NMR (500 MHz, TFA): δ 4.04 (3 H, s), 4.08 (3 H, s), 7.15 (2 H, d), 7.25 (2 H, d), 8.19 (2 H, d), 8.30 (2 H, d), 9.27 (1 H, s) and
15 9.37 (1 H, s) ppm.

Example 107 - synthesis of a mixture of 2-amino-6-(p-methoxyphenyl)pterin and 2-amino-7-(p-methoxyphenyl)pterin

The mixture obtained in example 106 (1.2 g, 4.5 mmoles) was
20 suspended in NaOH 1 N (80 ml) and refluxed till a solution was obtained. The hot solution was treated with acetic acid till pH 5, then cooled down and the resulting precipitate was filtered off and washed with water, ethanol and diethyl ether, providing a mixture of 2-amino-6-(p-methoxyphenyl)pterin and 2-amino-7-(p-methoxyphenyl) pterin as a yellow powder (1 g, yield: 82 %). Mass
25 spectrum: 270 ([M+H]⁺, 100).

Example 108 - synthesis of 2-acetylamino-6-(p-methoxyphenyl)pterin and 2-acetylamino-7-(p-methoxyphenyl)pterin

A suspension of the mixture obtained in example 107 (7.43 mmoles, 2
30 g) was suspended in a mixture of acetic anhydride (50 ml) and acetic acid (50 ml). The suspension was refluxed for 4 hours till a clear solution was obtained. Some insoluble material was filtered off and the solution was partly evaporated till precipitation starts. Further precipitation was achieved overnight

in the refrigerator. The resulting precipitate was filtered off and washed with ethyl acetate and diethyl ether, providing a mixture of 2-acetylamino-6-(p-methoxyphenyl)pterin and 2-acetylamino-7-(p-methoxyphenyl)pterin as a yellow powder (2.1 g, 91 % yield). Mass spectrum: 312 ($[M+H]^+$, 100).

5

Example 109 - synthesis of 2-acetylamino-4-(1,2,4-triazolyl)-6-(p-methoxyphenyl) pteridine and 2-acetylamino-4-(1,2,4-triazolyl)-7-(p-methoxyphenyl) pteridine

To a suspension of the mixture obtained in example 108 (1.5 g, 4 mmols) in dry pyridine (100 ml) was added 1,2,4-triazole (830 mg, 12 mmols) and 4-chlorophenyl phosphorodichloridate (1 ml, 6 mmols). The suspension was stirred for 2 days at room temperature under nitrogen. The solvents were removed *in vacuo*. The solid material was suspended in dichloromethane and washed with 2 % HCl. Evaporation of the solvents provided a mixture of 2-acetylamino-4-(1,2,4-triazolyl)-6-(p-methoxyphenyl) pteridine and 2-acetylamino-4-(1,2,4-triazolyl)-7-(p-methoxyphenyl) pteridine.

15

Example 110 - synthesis of 2-amino-4-isopropoxy-7-(p-methoxyphenyl) pteridine

To a suspension of the mixture obtained in example 109 (180 mg, 0.50 mmole) in isopropanol (8 ml) was added sodium (23 mg, 1 mmole). The suspension was stirred at room temperature overnight. The solvents were evaporated and the residue was purified by preparative TLC (silica, using a methanol/ CH_2Cl_2 (7:93) mixture as the eluent). At this stage, both regioisomers obtained were separated, thus providing the pure title compound as a yellow powder (yield: 45 %) which was further characterized by its mass spectrum: 312 ($[M+H]^+$, 65), 270 ($[M+H-propene]^+$, 100).

20

25

Example 111 - synthesis of 2-amino-4-isopropoxy-7-(3,4-dimethoxyphenyl) pteridine

30

The sequence of reactions described in examples 106 to 110 was followed, however starting from 3,4-dimethoxyphenylglyoxal instead of 4-methoxyphenylglyoxal in the first step. This provided 2-amino-4-isopropoxy-7-

(3,4-dimethoxyphenyl) pteridine, a compound which was further characterized by its mass spectrum: 342 ($[M+H]^+$, 55), 300 ($[M+H\text{-propene}]^+$, 75).

Example 112 - synthesis of 2-amino-4-ethoxy-7-(3,4-dimethoxyphenyl)

5 pteridine

The sequence of reactions described in examples 106 to 110 was followed, however starting from 3,4-dimethoxyphenylglyoxal instead of 4-methoxyphenylglyoxal in the first step, and from ethanol instead of isopropanol in the last step. This provided 2-amino-4-ethoxy-7-(3,4-dimethoxyphenyl)

10 pteridine, a compound which was further characterized as follows:

- MS: 328 ($[M+H]^+$, 100), 300 ($[M+H\text{-ethene}]^+$, 40);
- $^1\text{H-NMR}$ (500 MHz, $\text{DMSO-}d_6$): δ 1.44 (3 H, t), 3.86 (3 H, s), 3.88 (3 H, s), 4.54 (2 H, q), 7.13 (1 H, d), 7.16 (2 H, br s), 7.85 (1 H, d), 7.88 (1 H, dd) and 9.06 (1 H, s) ppm;
- 15 - $^{13}\text{C-NMR}$ (125 MHz, $\text{DMSO-}d_6$): δ 14.25, 55.67, 55.76, 63.06, 110.39, 111.89, 121.13, 121.25, 128.24, 136.87, 149.28, 151.62, 155.82, 156.72, 162.03 and 166.70 ppm.

Example 113 - synthesis of 2-amino-4-methoxy-7-(3,4-dimethoxyphenyl)

20 pteridine

The sequence of reactions described in examples 106 to 110 was followed, however starting from 3,4-dimethoxyphenylglyoxal instead of 4-methoxyphenylglyoxal in the first step, and from methanol instead of isopropanol in the last step. This provided 2-amino-4-ethoxy-7-(3,4-dimethoxyphenyl) pteridine, a compound which was further characterized by its mass spectrum: 314 ($[M+H]^+$, 100), 300 ($[M+H\text{-methane}]^+$, 20).

Example 114 – lymphocyte activation tests

Pteridine derivatives were first dissolved (10mM) in dimethylsulfoxide (hereinafter referred as DMSO) and further diluted in culture medium before use for the following *in vitro* experiments. The commercially available culture medium consisted of RPMI-1640 + 10% foetal calf serum (FCS). Some

pteridine derivatives described the previous examples were tested in the following lymphocyte activation tests:

Mixed Lymphocyte Reaction

Peripheral blood mononuclear cells (hereinafter referred as PBMC) were isolated from heparinized peripheral blood by density gradient centrifugation over Lymphoprep (Nycomed, Maorstua, Norway). Allogeneic PBMC or Epstein-Barr Virus-transformed human B cells [commercially available under the trade name RPMI1788 (ATCC name CCL156)] which strongly express B7-1 and B7-2 antigens were used as stimulator cells after irradiation with 30 Gy. MLR was performed in triplicate wells. After 5 days incubation at 37°C, 1 µCi [³H]-thymidine was added to each cup. After a further 16 hours incubation, cells were harvested and counted in a β-counter. Inhibition of proliferation by a compound (drug) described in some of the previous examples was counted using the formula:

$$\% \text{ inhibition} = \frac{(\text{cpm} + \text{drugs}) - (\text{cpm cult. med})}{(\text{cpm} - \text{drugs}) - (\text{OD cult. med})} \times 100$$

wherein cpm is the thymidine count per minute. The MLR assay is regarded by those skilled in the art as an *in vitro* analogue of the transplant rejection since it is based on the recognition of allogeneic major histocompatibility antigens on the stimulator leukocytes, by responding lymphocytes.

Assays for CD3 and CD 28

T cells were purified by removing non-T cells. Briefly, monocytes were removed by cold agglutination. The resulting lymphoid cells were further purified by a cell enrichment immunocolumn [Celect Human T commercially available from Biotex, Edmonton, Alberta, Canada]] by a process of negative selection. More than 95% of the B cells were removed with this procedure. After depletion, the resulting T cell preparation was highly purified, i.e. these cells could not be activated by phytohaemagglutinin (PHA) or rIL-2 alone at concentrations capable of stimulating RBMC prior to deletion.

Highly purified T cells (10⁶/ml) were stimulated by immobilized anti-CD3 or anti-CD28 monoclonal antibodies (hereinafter referred as mAb) in the presence of phorbol myristate acetate (hereinafter referred as PMA). Anti-CD3

mAb (available from CLB, Amsterdam, Netherlands) were fixed on a 96-microwell plates by incubating the wells with 50 µl of mAb solution (1/800 dilution in culture medium). For anti-CD28 mAb (available from CLB, Amsterdam, Netherlands) 50 µl (1/650 dilution in culture medium) was added directly to the wells. Further, a 20 µl PMA (commercially available from Sigma, St. Louis, Missouri, USA) solution (final concentration: 0.5 ng/ml) was added. Subsequently, 20 µl of a pteridine derivative described in the previous examples were added by serial dilution in triplicate wells. Finally 100 µl of the T-cell suspension (10^6 /ml) was added. After 48-hour incubation at 37°C in 5% CO₂, 20 µl of a bromo-deuridine (hereinafter referred as BrdU) 100 µM solution (commercially available as Cell Proliferation Elisa from Boehringer-Mannheim Belgium) was added to each well. After a further overnight incubation, T-cell proliferation was measured using a colorimetric immunoassay for qualification of cell proliferation based on the incorporation of BrdU during DNA synthesis. Optical density (hereinafter referred as OD) was measured by a Behring EL311 plate reader at 450 nm (reference wavelength: 690 nm). Inhibition of proliferation by the pteridine derivative (drug) was counted while using the formula:

$$\% \text{ inhibition} = \frac{(\text{OD} + \text{drugs}) - (\text{OD cult. med})}{(\text{OD} - \text{drugs}) - (\text{OD cult. med})} \times 100$$

Table 1 (wherein ND means not determined) below shows the IC₅₀ values for various pteridine derivatives in the MLR test and in the CD3 and CD28 assays. The IC₅₀ value represents the lowest concentration of the pteridine derivative (expressed in µmole/l) that resulted in a 50% suppression of the MLR or a 50% reduction in T-cell proliferation (for the CD3 and CD28 assay).

TABLE 1

Example n°	MLR	CD 3	CD28
1	15	110	80
2	12	4.0	12
3	25	160	100
4	> 200	140	110
5	140	> 200	> 200

TABLE 1 (follow)

Example n°	MLR	CD 3	CD28
6	5.0	15	15
7	3.6	4.5	3.9
8	12	7.5	7.5
9	15	80	20
10	17	50	20
11	12	20	20
12	0.08	0.5	0.1
13	75	25	25
14	100	75	20
15	50	20	15
16	3.5	4.3	3.2
17	15	25	25
18	13	9.7	6.7
19	8.5	5.7	2.8
20	2.2	0.8	0.7
21	15	100	20
22	12	20	20
23	15	125	20
24	0.1	0.4	0.3
25	15	80	20
26	15	70	25
27	4.6	14	12
28	0.4	0.5	0.4
29	15	40	25
30	4.8	11	4.2
31	12.3	11.7	4.2
32	2.0	7.3	6.5
33	22	17	3.7
34	12	11.7	5.0
35	2.0	20	15

TABLE 1 (follow)

Example n°	MLR	CD 3	CD28
36	10.5	14	4.5
37	2.8	14.8	4.5
38	++	++	++
39	++	+	+
40	++	++	++
41	0	+	+
42	+++	++	++
43	++	++	++
44	++	+	+
45	0	+	+
46	++	++	+
47	+++	++	++
48	+	+	+
49	++	++	++
50	++	+	+
51	+++	ND	ND
52	++	+	+
53	ND	+++	+++
68	8.3	20	7.0
69	> 10	9.0	4.0
71	10	10	0.8
73	ND	15	0.7
78	8.2	10	10
83	0.9	0.7	0.7
85	10	ND	ND
98	0.4	6.7	1.7

0 stands for concentrations of at least 151 μM ,

+

 stands for concentrations 16-150 μM ,

++ stands for concentrations 1-15 μM , and

+++ stands for concentrations lower than 1 μM .

The above data show that, whereas known immunosuppressant drugs like CyA are known to be active only in the CD3 assay, the pteridine derivatives according to the present invention were also active in the CD28 assay which is Ca^{2+} -calmodulin resistant. The CD28 pathway is a so-called cosignal pathway which is important for inducing energy and even tolerance in T-cells.

Example 115 - TNF- α and IL-1 β assays

Peripheral blood mononuclear cells (herein referred as PBMC), in response to stimulation by lipopolysaccharide (LPS), a gram-negative bacterial endotoxin, produce various chemokines, in particular human TNF- α and IL-1 β . Inhibition of the activation of PBMC can therefore be measured by the level of suppression of the production of TNF- α or IL-1 β by PBMC in response to stimulation by LPS.

Such inhibition measurement was performed as follows: PBMC were isolated from heparinized peripheral blood by density gradient centrifugation over Lymphoprep (commercially available from Nycomed, Norway). LPS was then added to the PMBC suspension in complete medium (10^6 cells /ml) at a final concentration of 1 $\mu\text{g/ml}$. The pteridine derivative to be tested was added at different concentrations (0.1 μM , 1 μM and 10 μM) and the cells were incubated at 37°C for 72 hours in 5% CO_2 . The supernatants were collected, then TNF- α and/or IL-1 β concentrations were measured with respectively an anti-TNF- α antibody or an anti-IL-1 β antibody in a sandwich ELISA (Duo Set ELISA human TNF α , commercially available from R&D Systems, United Kingdom). The colorimetric reading of the ELISA was measured by a Multiskan RC plate reader (commercially available from ThermoLabsystems, Finland) at 450 nm (reference wavelength: 690 nm). Data analysis was performed with Ascent software 2.6. (also from ThermoLabsystems, Finland): a standard curve (recombinant human TNF α) was drawn and the amount (pg/ml) of each sample on the standard curve was determined.

The % suppression of human TNF α production or human IL-1 β by the pteridine derivatives of the invention (drugs) was calculated using the formula:

$$\% \text{ suppression} = \frac{\text{pg/ml in drugs} - \text{pg/ml in cult. med.}}{(\text{pg/ml in cult. med.} + \text{LPS}) - \text{pg/ml cult. med.}}$$

Table 2 below shows the IC₅₀ values (expressed in µM) of the tested pteridine derivatives in the TNF-α assay.

5

TABLE 2

Example n°	TNF-α	Example n°	TNF-α
59	8.0	83	6.6
63	4.0	85	6.7
65	9.1	87	7.5
66	4.5	89	9.1
67	10.0	90	6.6
68	10.0	91	6.2
72	8.5	92	10.0
77	8.1	94	7.6
78	8.5	95	6.3
80	6.8	97	9.1
81	10.0	98	3.5

Example 116 – inhibition of the release of cytokines involved in asthma

The pteridine derivative of example 24 was tested, according to experimental procedures well established in the art, in the inhibition of two cytokines, IL-5 and IL-10, known to be involved in the development of asthma and other forms of severe allergic conditions. At the concentration of 0.1 µM/l, i.e. the IC₅₀ value of this derivative in the MLR test, the said derivative inhibited the release of IL-5 and IL-10 by 70% and 39%, respectively.

15 Example 117 – inhibition of the metastasis of melanoma cells in mice

C57BL/6 mice were injected with 1.5×10^6 B16BL/6 melanoma cells and were divided into 2 groups. Five days after the injection of the tumor cells, the two groups of animals were administered intraperitoneously, three times a week for two subsequent weeks, either (treated group) 20 mg/kg of the pteridine derivative of example 24 or (control group) the vehicle (5% DMSO in phosphate buffered saline). At the end of the experiment, all surviving tumor-bearing mice were sacrificed and examined in order to assess the absence or presence of the black metastases in inguinal and/or para-aortic lymph nodes (macroscopic metastasis). The findings were confirmed by histology, and the experiment done twice. The pooled results of the two experiments show that 80% of animals from the control group had metastases whereas only 33% of animals from the treated group had metastases.

Example 118 - *in vivo* leukocyte activation and immunosuppression whole blood assay

In response to non-self antigens (e.g. microbial antigens, allo-antigens, altered self antigens and auto-antigens), immune cells such as lymphocytes, NK cells, monocytes/macrophages and dendritic cells are activated to generate antigen-reactive effector cells. Activated immune cells up-regulate a number of pro-inflammatory genes encoding cytokines (e.g. IL-2, IFN- γ , and TNF- α), chemokines (e.g. monocyte chemo-attractant protein-1 (MCP-1), macrophage inflammatory protein-1 (MIP-1) and RANTES), and cell surface molecules. These leukocyte surface molecules may include activation associated molecules (e.g. CD69, the lymphocyte early activation molecule, and CD71, the transferrin receptor), adhesion molecules (e.g. ICAM-1, LFA-1 and L-selectins), co-stimulatory molecules (e.g. CD28, B7-1 and CD40L) and receptors for cytokines and chemokines (e.g. IL-2R and CCR1).

Immunomodulatory therapies are active by interfering with the responses of the immune system. Reliable, easy, fast and quantitative methods for monitoring the evolution of the leucocyte surface and cytoplasmic phenotypes

during activation are critical in evaluating the functional competence of the immune cells (e.g., during HIV infection), as well as the efficacy of immunoregulatory therapies in order to achieve optimal therapeutic effects and minimal side effects. Moreover, examination of drug action on leukocyte
5 activation is able to provide critical evaluation of pharmacokinetic and pharmacodynamic effects of immunomodulatory agents. Within this context, the following provides a useful *in vivo* leukocyte activation and immunosuppression whole blood assay (hereinafter referred as WBA).

Inbred mice (e.g. AKR strain), weighting from 20 to 30g, were stimulated
10 by intra-peritoneal injection with mitogens specific for T cells (e.g. Concanavalin A, hereinafter referred as ConA, 500 µg per mouse), B cells or monocytes/macrophages (e.g. LPS 20 µg per mouse). The pteridine derivatives of example 83 and, respectively, example 98, were administered intraperitoneously (30 mg/kg per day, administered 2 hours before ConA injection).
15 Peripheral blood was collected 24, 36, 48, and respectively 72 hours after stimulation for immune fluorescence staining and FACScan analysis.

FACScan analysis proceeded as follows: heparinized whole blood (50 µl) was removed of red blood cells by incubation with a lysing buffer. The remaining leukocytes were double-stained with phycoerythrin-conjugated antibodies specific
20 for T cells (e.g. T cell receptor TCR), B cells (e.g. CD45R/B220 (commercially available from Pharmingen) or monocytes/macrophages (e.g. F4/80) (commercially available from Serotec) in combination with FITC-conjugated antibodies against various cell surface markers such as CD69 (early activation molecule), CD25 (IL-2 receptor), CD71 (transferring receptor), B7-1/2 (ligand for the co-
25 stimulatory molecule CD28) and CD11b (complement receptor 3). Cells were examined by flow cytometry using CellQuest software (commercially available from Decton Dickinson).

Peripheral blood T cells expressed an up-regulated level of CD69, representing up to 3% of TCR⁺,CD69⁺ cells respectively 1 and 2 days following
30 ConA injection, as compared to baseline levels (0.5%). T cells from mice treated

with the pteridine derivatives of examples 83 and 98 show a remarkable reduction of this response to 1.6% and 2.4% respectively.

Peripheral blood T cells showed a rapid up-regulation of CD71 from baseline level (4.8%) to 9.3% and 5.5% after 1 day and 2 days, respectively. The
5 pteridine derivatives of examples 83 and 98 suppressed ConA-stimulated CD71 up-regulation on T cells to baseline levels.

Peripheral blood T cells expressed increased levels of IL-2R with up to 3% of TCR+, IL-2R+ cells 1 and 2 days following ConA stimulation, with peak levels after day 2. Administration of the pteridine derivatives of examples 83 and 98
10 effectively suppresses this effect by reducing expression to 1.5% and 1.7%, respectively.

Up to 6.6 % of peripheral blood T cells displayed an up-regulation of the CD134 molecule expression within 2 days following ConA stimulation, i.e. a two-fold increase of baseline levels (2.2%). Treatment with the pteridine derivatives of
15 examples 83 and 98 suppressed this response to 3.5% and 4.2%, respectively.

This FACScan-based cell surface phenotype analysis for *in vivo* leukocyte activation, namely for an *in vivo* whole blood assay, demonstrates the immunosuppressive effects of the pteridine derivatives of examples 83 and 98 *in vivo*. This observation is consistent with the above experimental results *in vitro*
20 showing that both compounds inhibit T cell activation and proliferation in mixed lymphocyte culture, and in the CD3 and CD28 assays.

In vivo WBA by injecting immune activators, followed by FACS analysis of leukocyte surface phenotypes in the whole blood can be applied to monitor *in vivo* activation of different types of leukocytes such as T cells, B cells, monocytes/macrophages, and dendritic cells. This technology allows simple,
25 rapid, direct, stable and quantitative evaluation of the immune competence of the said leukocytes, as well as pharmacokinetic and pharmacodynamic profiles of the pteridine derivatives used as immunomodulatory agents *in vivo*.

Despite various *in vitro* models are used for the discovery of new immunosuppressive drugs, there are essentially no ideal animal models that are simple, fast, and cost-effective. Models of organ transplantation (e.g. hearts, kidney and skin) in rodents are limited by technical difficulty, and cannot be used for primary screening of a large numbers of candidate drugs. Although recently a mouse model of drug-mediated immunosuppression based on rejection of an allogeneic subcutaneous tumor was reported, several significant limitations should be considered. For example first, the host immune responses against this specific tumor seems to be T cell-dependent, and thus the activity of a drug on other cell types such as B cells, monocytes/macrophages and dendritic cells cannot be evaluated by using this model. Secondly, immunosuppressive drugs with anti-proliferation activity (e.g. rapamycin) may play a direct role on tumor angiogenesis and tumor cell growth. Thirdly, the feature of chronic rejection of the tumor (e.g. after 14 days) prevents from evaluating the pharmacodynamic properties of drugs. Finally, this animal tumor rejection model is inaccurate (e.g. quantified by size measurement) and time-costly (at least 14 days).

FACScan-based *in vitro* WBA has been used to investigate the pharmacodynamic effects of immunomodulatory drugs administrated *in vivo*, but only by taking blood from treated recipients and stimulating their lymphocytes *in vitro*. These methods show remarkable advantages as compared to the purified peripheral blood mononuclear cell (PBMC) cultures in terms of maintaining cell viability, as well as the effects within the whole blood of immunomodulatory drugs previously administrated *in vivo*. However, *in vitro* WBA has important flaws. First, the evaluation of the drug effects uses an *in vitro* activation system, including manipulation, dilution and stimulation of the blood, and subsequent culture of the blood in the presence of an exogenous serum (e.g. foetal calf serum). This *in vitro* activation system may not necessarily and accurately reflect the effects of drugs on leukocyte activation *in vivo*, nor does this assay give information on the absorption or metabolite production of drugs that may occur *in vivo*. Secondly, *in vitro* stimulation of the whole blood by mitogens (e.g. Con A, PHA or LPS) causes aggregation of activated leukocytes, platelets and red blood


cells, as well as generation of spontaneous fluorescence by *in vitro* damaged cells. These alterations frequently interfere with the FACS analysis. Thirdly, plastic adhesion of monocytes/macrophages during *in vitro* culture may induce additional steps in order to examine these cell types. Finally, *in vitro* culture of leucocytes is expensive, time consuming, and poorly reproducible. The above described *in vivo* WBA assay is able to overcome the above mentioned problems and may be characterized as follows:

- leukocyte stimulation is performed *in vivo* (e.g. by *in vivo* injection of leucocyte stimulating agents such as, but not limited to, ConA or LPS, thus enabling the full *in vivo* evaluation of the effect of drugs on leucocyte activation in physiological conditions,
- various stimuli can be used that are specific for different cell types, including T cells (e.g. ConA, anti-CD3 antibodies and alloantigens), B cells (e.g. T cell-independent xeno-antigens, sheep red blood cells, TNP-ficoll, TNP-BA, anti-IgM antibodies and LPS) and monocytes/macrophages (e.g. LPS, TNF- α and IFN- γ). Given the time-dependent nature of the expression of cell surface antigens, the effects of drugs can be studied on early as well as on late leucocyte activation, e.g. by analyzing the expression of early (e.g. CD69 and CD71) or late (e.g. IL-2R and CD134) appearing antigens,
- by varying the time of administration of a drug prior to or after *in vivo* injection of the leucocyte stimuli, the pharmacodynamic profile of the said drug can be determined. Similarly, the effects and the duration of activity of a drug via different routes of administration, e.g. intraperitoneously, intramuscularly and orally, can be studied,
- by using the same test in immunodeficient animals (e.g. T cell-deficient nude mice), the effect of a drug on selected immune phenomena can be investigated (e.g. on *in vivo* T cell-independent B cell activation in nude mice; T cell activation in T cell-reconstituted SCID mice that have congenital defect on both T and B cells),

- peripheral leukocytes from different sources, such as spleen, lymph nodes and peritoneal cavity, can be tested,
- in combining FACScan-based intracellular phenotype analysis with e.g. intracellular cytokine staining, or DNA synthesis by BrdU or 5- and 6-carboxyfluorescein diacetate succinimidyl ester (CFSE) incorporation, or cell cycle studies, the information obtained can be further broadened, and
- the method is able to help in elucidating the mechanism underlying *in vivo* up-regulation of expression of certain activation molecules, such as CD69, CD25, CD71, CD134 and B7-1/2, in various types of peripheral blood leucocytes.

CLAIMS

1. A pteridine derivative having the general formula:



wherein X represents an oxygen atom or a group with the formula $S(O)_m$ wherein m

5 is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R_1 is a group selected from the group consisting of C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-7} alkenyl, C_{2-7} alkynyl, halo C_{1-4} alkyl, C_{3-10} cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C_{1-7} alkyl, thio C_{3-10} cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenyl-amino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenyl-hydrazino; or R_1 is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;
- Z is a group independently defined as R_1 or Z is hydrogen or the group NZ together with R_1 is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R_2 is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl, ureido; thio-ureido, sulfonamido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino;

- alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C₃₋₇ alkylamino; arylamino; arylalkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; thio C₁₋₇ alkyl; 5 arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C₃₋₁₀ cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals;
- R₄ is an atom or a group selected from the group consisting of hydrogen; halogen; C₁₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; carboxy C₁₋₇ 10 alkyl; carboxyaryl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₁₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-acetal; carboxylic acid; carboxylic 15 acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido; alkylamino; cycloalkyl-amino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercaptoalkylamino; heterocyclic 20 amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and thioamides thereof; phenyl substituted with one or more substituents selected from the group consisting of C₁₋₇ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, 25 hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, 30 sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino, thioalkyl-amino, acylamino, thioacylamino, cyano, carboxylic acid or esters or

thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynyl-amino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, 5 heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; aryl groups other than phenyl, the said aryl groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, 10 oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino, thioalkyl-amino, acylamino, thioacylamino, cyano, carboxylic acid or esters or 15 thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynyl-amino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; 20 optionally substituted heterocyclic radicals selected from the group consisting of oxabicycloheptyl, azabenzimidazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl, tetrahydrofuryl, tetrahydro-pyranyl, tetrahydropyranyl, tetrahydroquinoleinyl, tetrahydro-thienyl and dioxide thereof, dihydrothienyl dioxide, dioxindolyl, dioxinyl, dioxenyl, 25 dioxazinyl, thioxanyl, thioxolyl, thio-urazolyl, thiotriazolyl, thiopyranyl, thiopyranyl, coumarinyl, quinoleinyl, oxyquinoleinyl, quinuclidinyl, xanthi-nyl, dihydropyranyl, benzodihydrofuryl, benzothiopyranyl, benzothio-pyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl, benzodioxanyl, benzothiadiazolyl, benzotriazinyl, benzothiazolyl, benzoxazolyl, 30 phenothioxinyl, phenothiazolyl, phenothieryl, phenopyranyl, phenoxazolyl, pyridinyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl, thiomorpholinyl,

pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl, triazolyl,
 benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl,
 isothiazolyl, oxazolyl, oxadiazolyl, pyrrolyl, furyl, dihydrofuryl, furoyl,
 hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithiinyl, thienyl,
 5 indolyl, indazolyl, benzofuryl, quinolyl, quinazolinyl, quinoxalyl,
 carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothienyl,
 naphthothienyl, thianthrenyl, pyranyl, pyronyl, benzopyronyl, isobenzofuranyl,
 chromenyl, phenoxathiinyl, indoliziny, quinoliziny, isoquinolyl,
 phthalazinyl, naphthiridinyl, cinnoliny, pteridinyl, carboliny, acridiny,
 10 perimidiny, phenanthroliny, phenazinyl, phenothiazinyl, imidazoliny,
 imidazolidiny, benzimidazolyl, pyrazoliny, pyrazolidiny, pyrroliny,
 pyrrolidinyl, piperazinyl, uridiny, thymidinyl, cytidiny, aziriny, aziridinyl,
 diaziriny, diaziridinyl, oxirany, oxaziridinyl, dioxirany, thiirany, azetyl,
 dihydroazetyl, azetidiny, oxetyl, oxetany, thietyl, thietany, diazabicyclo-
 15 octyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromany, chromanonyl,
 thiochromany, thiochromanonyl, thiochromenyl, benzofurany, benzisothiazolyl,
 benzocarbazolyl, benzochromonyl, benzisoalloxazinyl, benzocoumariny, thiocoumariny,
 phenometoxazinyl, phenoparoxazinyl, phentriazinyl, thiodiazinyl, thiodiazolyl,
 20 indoxyl, thioindoxyl, benzodiazinyl, phtalidyl, phtalimidiny, phtalazonyl, alloxazinyl,
 xanthionyl, isatyl, isopyrazolyl, isopyrazolonyl, urazolyl, uraziny, uretiny, uretidiny,
 succiny, succinimido, benzylsultimyl and benzylsultamyl; aromatic or heterocyclic
 substituents substituted with an aliphatic spacer between the pteridine ring
 and the aromatic or heterocyclic substituent, whereby said aliphatic spacer
 25 is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4
 carbon atoms which may contain one or more functions, atoms or radicals
 selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol
 (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino,
 oximino, alkylloximino, amino-acid, cyano, acylamino, thioacylamino,
 30 carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or
 thioester or halide or anhydride or amide, thiocarboxylic acid or ester or

- thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 1 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, aminoacid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; and
- R₃ is an atom or a group selected from the group consisting of fluoro, bromo, iodo, C₂₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₂₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino; acylamino; thio-acylamino; thio-acetal; carboxylic acid; carboxylic acid esters, thioesters, amides, halides, anhydrides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, amides, halides, anhydrides and thioamides; hydroxyl; sulfhydryl; nitro; carbamoyl; thiocarbamoyl; ureido; thio-ureido; amino; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercaptoalkylamino; heterocyclic amino; heterocyclic-substituted

alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenyl-
 hydrazino; cysteinyl acid, esters, thioesters, amides and thioamides
 thereof; aryl optionally substituted with one or more substituents selected
 from the group consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl,
 5 C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀
 cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted
 alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic,
 arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl,
 thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, mercapto-
 10 amino, alkoxyamino, thioalkylamino, acylamino, thio-acylamino, cyano,
 carboxylic acid or esters or thioesters or halides or anhydrides or amides
 thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides
 or amides thereof, alkylamino, cycloalkylamino, alkenylamino,
 cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxy-
 15 alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino,
 alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic
 radicals; aromatic or heterocyclic substituents substituted with an aliphatic
 spacer between the pteridine ring and the aromatic or heterocyclic
 substituent, whereby said aliphatic spacer is a branched or straight,
 20 saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may
 contain one or more functions, atoms or radicals selected from the group
 consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether,
 thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-
 acid, cyano, carboxylic acid or ester or thioester or amide, nitro, thio C₁₋₇
 25 alkyl, thio C₃₋₁₀ cycloalkyl, alkylamino, cycloalkylamino, alkenylamino,
 cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxy-
 alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkyl-
 hydrazino, phenylhydrazino, sulfonyl, sulfonamido and halogen; branched
 or straight, saturated or unsaturated aliphatic chains of 2 to 7 carbon
 30 atoms optionally containing one or more functions selected from the group
 consisting of thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, thio-

acetal, amino, imino, oximino, alkylloximino, amino-acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thio carboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, aryl-amino, arylalkylamino, hydroxy-alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; or R₃ together with R₄ forms a homocyclic or heterocyclic radical;

and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-*N*-oxide thereof and/or a solvate thereof and/or a dihydro- or tetrahydropteridine derivative thereof.

2. A pteridine derivative according to claim 1, wherein R₁ is selected from the group consisting of methyl, ethyl, isopropyl and pentyl.

3. A pteridine derivative according to claim 1, wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents.

4. A pteridine derivative according to claim 1, wherein R₃ is a phenyl group with one or more substituents each independently selected from the group consisting of fluoro, methoxy, ethoxy, trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate.

5. A pteridine derivative according to claim 1, wherein:

- X is NZ,
- Z is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl and benzyl, and

- R₁ is selected from the group consisting of methyl, ethyl, n-propyl and benzyl.

6. A pteridine derivative according to claim 1, wherein X is NZ and wherein the
 5 group NZ together with R₁ is selected from the group consisting of hydroxylamino, morpholinyl, piperidinyl, piperazinyl, 1,2,4-triazolyl and N-methylpiperazinyl.

7. A pteridine derivative according to claim 1, being selected from the group
 10 consisting of:


- 2-amino-4-ethoxy-6-(4-methoxyphenyl)-pteridine
- 2-amino-4-ethoxy-6-(2-methoxyphenyl)-pteridine
- 2-amino-4-ethoxy-6-(3-methoxyphenyl)-pteridine
- 2-amino-4-ethoxy-6-(3,4-difluorophenyl)-pteridine
- 15 - 2-amino-4-ethoxy-6-(4-dimethylaminophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-trifluoromethylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(2-thienyl)-pteridine
- 2-amino-4-ethoxy-6-(3-thienyl)-pteridine
- 2-amino-4-ethoxy-6-(3,4-dichlorophenyl)-pteridine
- 20 - 2-amino-4-ethoxy-6-(4-cyanophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-ethoxyphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-fluorophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-ethylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-acetylphenyl)-pteridine
- 25 - 2-amino-4-ethoxy-6-(3-fluoro-4-methylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-thiomethylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(3-methyl-4-methoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethylphenyl)-pteridine
- 30 - 2-amino-4-isopropoxy-6-(3-chloro-4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3-chloro-4-fluorophenyl)-pteridine

- 2-amino-4-isopropoxy-6-(4-N,N-diethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-difluorophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-methoxyphenyl)-pteridine
- 5 - 2-amino-4-isopropoxy-6-(4-ethoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(3-thienyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-cyanophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-benzoic acid methyl ester)-pteridine
- 10 - 2-amino-4-isopropoxy-6-(4-acetylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-ethylthio-6-(3,4-dimethoxyphenyl)-pteridine,
- 2-amino-4-isopropylthio-6-(3,4-dimethoxyphenyl)-pteridine,
- 2-amino-4-pentoxy-6-styrylpteridine,
- 15 - 2-amino-4-n-pentoxy-6-(1,2-dibromo-2-phenylethyl)-pteridine,
- 2-amino-4-methoxy-6-styryl-7-methoxypteridine,
- 2,4-diamino-6-phenyl-7-methylpteridine,
- 2-amino-4-dimethylamino-6-phenylpteridine,
- 2-amino-4-dimethylamino-6-(4-tolyl)pteridine,
- 20 - 2-amino-4-dimethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-phenylpteridine,
- 2-amino-4-diethylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-diethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 25 - 2-amino-4-dibenzylamino-6-phenyl pteridine,
- 2-amino-4-dibenzylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dipropylamino-6-phenylpteridine,
- 30 - 2-amino-4-dipropylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-dipropylamino-6-(4-methoxyphenyl)pteridine,

- 2-amino-4-dipropylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-phenylpteridine,
- 2-amino-4-morpholino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-morpholino-6-(4-methoxyphenyl)pteridine,
- 5 - 2-amino-4-morpholino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-piperidino-6-phenylpteridine,
- 2-amino-4-piperidino-6-(4-chlorophenyl) pteridine,
- 2-amino-4-piperidino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperidino-6-(3,4-dimethoxyphenyl)pteridine,
- 10 - 2-amino-4-N-methylpiperazino-6-phenylpteridine,
- 2-amino-4-N-methylpiperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-N-methylpiperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-methylpiperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-pyrrolidino-6-(4-methoxyphenyl)pteridine,
- 15 - 2-amino-4-piperazino-6-phenylpteridine,
- 2-amino-4-piperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-piperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(3,4,5-trimethoxyphenyl)pteridine,
- 20 - 2-amino-4-morpholino-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(3-adamantylamino)-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(3-adamantylamino)-6-naphtylpteridine,
- 2-amino-4-(4-adamantylamino)-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(4-adamantylamino)-6-naphtylpteridine,
- 25 - 2-amino-4-morpholino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-formylidene-3,4-dihydroxyphenyl)
pteridine,
- 2-amino-4-pyrrolidino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 30 - 2-amino-4-dimethylamino-6-methylpteridine,
- 2-amino-4-ethoxy-6-phenylpteridine,

- 2-amino-4-propylamino-6-phenylpteridine,
- 2-amino-4-propylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-acetamido-4-hydroxy-6-(3,4-dimethoxyphenyl)pteridine,
- 2-acetamido-4-isopropoxy-6-(3,4-dimethoxyphenyl)pteridine, and
- 5 - 2-amino-4-ethoxy-6-(3,4-dimethoxyphenyl)pteridine.

8. A pharmaceutical composition comprising as an active principle at least one pteridine derivative having the general formula:



- 10 wherein X represents an oxygen atom or a group with the formula S(O)_m wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:
- R₁ is a group selected from the group consisting of C₁₋₇ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₄ alkyl, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenyl-amino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; or R₁ is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;

- Z is a group independently defined as R₁ or Z is hydrogen or the group NZ together with R₁ is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R₂ is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl, ureido; thio-ureido, sulfonamido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino; alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C₃₋₇ alkylamino; arylamino; arylalkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; thio C₁₋₇ alkyl; arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C₃₋₁₀ cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals,
- R₄ is an atom or a group selected from the group consisting of hydrogen; halogen; C₁₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; carboxy C₁₋₇ alkyl; acetoxyc C₁₋₇ alkyl; carboxyaryl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₁₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-acetal; carboxylic acid; carboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercapto-alkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and thioamides thereof; phenyl substituted with one or more substituents selected from the group consisting of C₁₋₇ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl,

halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino, thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; aryl groups other than phenyl, the said aryl groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₇ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino, thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic radicals selected from the group consisting of oxabicycloheptyl, azabenzimidazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydropyranyl, tetrahydroquinoleinyl, tetrahydro-thienyl and dioxide thereof, dihydrothienyl dioxide, dioxindolyl, dioxinyl, dioxenyl, dioxazinyl, thioxanyl, thioxolyl, thiourazolyl, thiotriazolyl, thiopyranyl,

thiopyranyl, coumarinyl, quinoleinyl, oxyquinoleinyl, quinuclidinyl,
 xanthinyl, dihydropyranyl, benzodihydrofuryl, benzothiopyranyl, benzothio-
 pyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl, benzodioxanyl,
 benzothiadiazolyl, benzotriazinyl, benzothiazolyl, benzoxazolyl, pheno-
 5 thioxinyl, phenothiazolyl, phenothienyl, phenopyranyl, phenoxazolyl,
 pyridinyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl, thiomorpholinyl,
 pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl, triazolyl,
 benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl,
 isothiazolyl, oxazolyl, oxadiazolyl, pyrrolyl, furyl, dihydrofuryl, furoyl,
 10 hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithiinyl, thienyl,
 indolyl, indazolyl, benzofuryl, quinolyl, quinazolinyl, quinoxalinyl,
 carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothienyl,
 naphthothienyl, thianthrenyl, pyranyl, pyronyl, benzopyranyl, isobenzofuranyl,
 chromenyl, phenoxathiinyl, indoliziny, quinoliziny, isoquinolyl,
 15 phthalazinyl, naphthiridinyl, cinnoliny, pteridinyl, carbolinyl, acridinyl,
 perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, imidazolinyl,
 imidazolidinyl, benzimidazolyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl,
 pyrrolidinyl, piperazinyl, uridinyl, thymidinyl, cytidinyl, azirinyl, aziridinyl,
 diazirinyl, diaziridinyl, oxiranyl, oxaziridinyl, dioxiranyl, thiiranyl, azetyl,
 20 dihydroazetyl, azetidiny, oxetyl, oxetanyl, thietyl, thietanyl, diazabicyclo-
 octyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromanyl, chromanonyl,
 thiochromanyl, thiochromanonyl, thiochromenyl, benzofuranyl,
 benzisothiazolyl, benzocarbazolyl, benzochromonyl, benzisoalloxazinyl,
 benzocoumarinyl, thiocoumarinyl, phenometoxazinyl, phenoparoxazinyl,
 25 phentriazinyl, thiodiazinyl, thiodiazolyl, indoxyl, thio-indoxyl, benzodiazinyl,
 phtalidyl, phtalimidinyl, phtalazonyl, alloxazinyl, xanthionyl, isatyl,
 isopyrazolyl, isopyrazolonyl, urazolyl, urazinyl, uretinyl, uretidinyl, succinyl,
 succinimido, benzylsultimyl and benzylsultamyl; aromatic or heterocyclic
 30 substituents substituted with an aliphatic spacer between the pteridine ring
 and the aromatic or heterocyclic substituent, whereby said aliphatic spacer
 is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4

- carbon atoms which may contain one or more functions, atoms or radicals selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 2 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, aminoacid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxy-alkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; and
- R₃ is an atom or a group defined as R₄ or R₃ is selected from the group consisting of morpholinyl, amino, hydrogen, methyl, thiomethyl and chloro; or R₃ together with R₄ forms a homocyclic or heterocyclic radical; and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-N-oxide thereof and/or a solvate and/or a dihydro- or tetrahydropteridine derivative thereof.

9. A pharmaceutical composition according to claim 8, wherein R₁ is selected from the group consisting of methyl, ethyl, isopropyl and pentyl.
10. A pharmaceutical composition according to claim 8, wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents.
11. A pharmaceutical composition according to claim 8, wherein R₃ is a phenyl group with one or more substituents each independently selected from the group consisting of fluoro, methoxy, ethoxy, trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate.
12. A pharmaceutical composition according to claim 8, wherein:
- X is NZ,
 - Z is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl and benzyl, and
 - R₁ is selected from the group consisting of methyl, ethyl, n-propyl and benzyl.
13. A pharmaceutical composition according to claim 8, wherein X is NZ and wherein the group NZ together with R₁ is selected from the group consisting of hydroxylamino, morpholinyl, piperidinyl, piperazinyl, 1,2,4-triazolyl and N-methylpiperazinyl.
14. A pharmaceutical composition according to claim 8, wherein the pteridine derivative is a compound selected from the group consisting of:
- 2-amino-4-ethoxy-6-(4-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(2-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(3-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(3,4-difluorophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-dimethylaminophenyl)-pteridine

- 2-amino-4-ethoxy-6-(4-trifluoromethylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(2-thienyl)-pteridine
- 2-amino-4-ethoxy-6-(3-thienyl)-pteridine
- 2-amino-4-ethoxy-6-(3,4-dichlorophenyl)-pteridine
- 5 - 2-amino-4-ethoxy-6-(4-cyanophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-ethoxyphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-fluorophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-ethylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-acetylphenyl)-pteridine
- 10 - 2-amino-4-ethoxy-6-(3-fluoro-4-methylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-methylthiophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(3-methyl-4-methoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethylphenyl)-pteridine
- 15 - 2-amino-4-isopropoxy-6-(3-chloro-4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3-chloro-4-fluorophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-N,N-diethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-difluorophenyl)-pteridine
- 20 - 2-amino-4-isopropoxy-6-(4-methoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-ethoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(3-thienyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-cyanophenyl)-pteridine
- 25 - 2-amino-4-isopropoxy-6-(4-benzoic acid methyl ester)-pteridine
- 2-amino-4-isopropoxy-6-(4-acetylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-ethylthio-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-isopropylthio-6-(3,4-dimethoxyphenyl)-pteridine
- 30 - 2-amino-4-pentoxy-6-styrylpteridine,
- 2-amino-4-n-pentoxy-6-(1,2-dibromo-2-phenylethyl)-pteridine,

- 2-amino-4-methoxy-6-styryl-7-methoxypteridine,
- 2,4-diamino-6-phenyl-7-methylpteridine,
- 2-amino-4-dimethylamino-6-phenylpteridine,
- 2-amino-4-dimethylamino-6-(4-tolyl)pteridine,
- 5 - 2-amino-4-dimethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-phenylpteridine,
- 2-amino-4-diethylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-diethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 10 - 2-amino-4-dibenzylamino-6-phenyl pteridine,
- 2-amino-4-dibenzylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dipropylamino-6-phenylpteridine,
- 15 - 2-amino-4-dipropylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-dipropylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-dipropylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-phenylpteridine,
- 2-amino-4-morpholino-6-(4-chlorophenyl)pteridine,
- 20 - 2-amino-4-morpholino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-piperidino-6-phenylpteridine,
- 2-amino-4-piperidino-6-(4-chlorophenyl) pteridine,
- 2-amino-4-piperidino-6-(4-methoxyphenyl)pteridine,
- 25 - 2-amino-4-piperidino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-N-methylpiperazino-6-phenylpteridine,
- 2-amino-4-N-methylpiperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-N-methylpiperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-methylpiperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 30 - 2-amino-4-pyrrolidino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperazino-6-phenylpteridine,

- 2-amino-4-piperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-piperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dibenzylamino-6-(3,4,5-trimethoxyphenyl)pteridine,
- 5 - 2-amino-4-morpholino-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(3-adamantylamino)-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(3-adamantylamino)-6-naphtylpteridine,
- 2-amino-4-(4-adamantylamino)-6-(3,4,5-trimethoxyphenyl)pteridine,
- 2-amino-4-(4-adamantylamino)-6-naphtylpteridine,
- 10 - 2-amino-4-morpholino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine,
- 2-amino-4-pyrrolidino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 15 - 2-amino-4-dimethylamino-6-methylpteridine,
- 2-amino-4-ethoxy-6-phenylpteridine,
- 2-amino-4-propylamino-6-phenylpteridine,
- 2-amino-4-propylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-acetamido-4-hydroxy-6-(3,4-dimethoxyphenyl)pteridine,
- 20 - 2-acetamido-4-isopropoxy-6-(3,4-dimethoxyphenyl)pteridine, and
- 2-amino-4-ethoxy-6-(3,4-dimethoxyphenyl)pteridine.

15. A pharmaceutical composition according to claim 8, further comprising one or more biologically-active drugs selected from the group consisting of immunosuppressant and/or immunomodulator drugs, antineoplastic drugs, and antiviral agents.

16. A pharmaceutical composition according to claim 8, further comprising one or more immunosuppressant drugs selected from the group consisting of cyclosporin A; substituted xanthines; pentoxifylline; daltroban, sirolimus, tacrolimus; rapamycin and derivatives thereof; leflunomide or an active

metabolite or an analog thereof; mycophenolic acid and salts thereof; adrenocortical steroids; azathioprine, brequinar; gusperimus; 6-mercaptopurine; mizoribine; chloroquine; hydroxychloroquine; monoclonal antibodies with immunosuppressive properties; etanercept; infliximab; and kineret.

5

17. A pharmaceutical composition according to claim 8, further comprising one or more immunomodulator drugs selected from the group consisting of acemannan, amiprilose, bucillamine, ditiocarb sodium, imiquimod, Inosine Pranobex, interferon- β , interferon- γ , lentinan, levamisole, pidotimod, romurtide, platonin,
10 procodazole, propagermanium, thymomodulin, thymopentin and ubenimex.


18. A pharmaceutical composition according to claim 8, further comprising one or more antineoplastic drugs selected from the group consisting of alkaloids, alkylating agents, alkyl sulfonates, aziridines, ethylenimines, methylmelamines,
15 nitrogen mustards, nitrosoureas, antibiotics, antimetabolites, folic acid analogs, purine analogs, pyrimidine analogs, enzymes, interferon and platinum complexes.

19. A pharmaceutical composition according to claim 8, further comprising one or
20 more antiviral agents selected from the group consisting of retroviral enzyme inhibitors, HIV-1 IN inhibitors, nucleoside reverse transcriptase inhibitors, zidovudine, lamivudine, didanosine, stavudine, zalcitabine, non-nucleoside reverse transcriptase inhibitors, nevirapine, delavirdine, foscarnet sodium, HIV-1 protease inhibitors, saquinavir, ritonavir, indinavir, nelfinavir, acyclovir, cidofovir,
25 cytarabine, edoxudine, famciclovir, floxuridine, ganciclovir, idoxuridine, penciclovir, sorivudine, trifluridine, valaciclovir, vidarabine, kethoxal, methisazone, moroxydine, podophyllotoxin, ribavirine, rimantadine, stallimycine, statolon, tromantadine and xenazoic acid.

20. A method for the prevention or treatment in a patient of a pathologic condition selected from the group consisting of:

- immune and auto-immune disorders,
- cardiovascular disorders,
- 5 - disorders of the central nervous system, and
- cell proliferative disorders,

comprising the administration to the patient of an effective amount of a pharmaceutical composition comprising as an active principle at least one pteridine derivative having the general formula:



wherein X represents an oxygen atom or a group with the formula $S(O)_m$ wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R_1 is a group selected from the group consisting of C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, aryl, alkylaryl, arylalkyl, heterocyclic, heterocyclic-substituted alkyl and alkyl-substituted heterocyclic, each of said groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{2-7} alkenyl, C_{2-7} alkynyl, halo C_{1-4} alkyl, C_{3-10} cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C_{1-7} alkyl, thio C_{3-10} cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, hydroxyl, sulfhydryl, nitro, hydroxylamino, mercaptoamino, cyano, carboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, thiocarboxylic acid or esters or thioesters or amides or thioamides or halides or anhydrides thereof, carbamoyl, thiocarbamoyl, ureido, thio-ureido, amino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenyl-amino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and

phenyl-hydrazino; or R_1 is a carboxyalkyl, carboxyaryl, thiocarboxyaryl or thiocarboxyalkyl group;

- Z is a group independently defined as R_1 or Z is hydrogen or the group NZ together with R_1 is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R_2 is selected from the group consisting of amino; acylamino; thioacylamino; carbamoyl; thiocarbamoyl, ureido; thio-ureido, sulfonamido; hydroxylamino; alkoxyamino; thioalkylamino; mercaptoamino, hydrazino; alkylhydrazino; phenylhydrazino; optionally substituted heterocyclic radicals; C_{3-7} alkylamino; arylamino; arylalkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; heterocyclic amino; hydroxyalkylamino; mercaptoalkylamino; C_{1-7} alkoxy; C_{3-10} cycloalkoxy; thio C_{1-7} alkyl; arylsulfoxide; arylsulfone; heterocyclic sulfoxide; heterocyclic sulfone; thio C_{3-10} cycloalkyl; aryloxy; arylthio; arylalkyloxy; arylalkylthio; oxyheterocyclic and thioheterocyclic radicals,
- R_4 is an atom or a group selected from the group consisting of hydrogen; halogen; C_{1-7} alkyl; C_{2-7} alkenyl; C_{2-7} alkynyl; halo C_{1-7} alkyl; carboxy C_{1-7} alkyl; acetoxyl C_{1-7} alkyl; carboxyaryl; C_{1-7} alkoxy; C_{3-10} cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C_{1-7} alkyl; thio C_{3-10} cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; hydroxylamino; mercapto-amino; acylamino; thio-acylamino; alkoxyamino; thioalkylamino; acetal; thio-acetal; carboxylic acid; carboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido; alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynylamino; arylamino; arylalkylamino; hydroxyalkylamino; mercapto-alkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and

thioamides thereof; phenyl substituted with one or more substituents selected from the group consisting of C₁₋₄ alkyl, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₄ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxyamino, mercaptoamino, thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenyl-amino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; aryl groups other than phenyl, the said aryl groups being optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₇ alkenyl, C₂₋₇ alkynyl, halo C₁₋₄ alkyl, nitro, hydroxyl, sulfhydryl, amino, C₃₋₁₀ cycloalkoxy, aryloxy, arylalkyloxy, oxyheterocyclic, heterocyclic-substituted alkyloxy, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, thioaryl, thioheterocyclic, arylalkylthio, heterocyclic-substituted alkylthio, formyl, carbamoyl, thiocarbamoyl, ureido, thio-ureido, sulfonamido, hydroxylamino, alkoxy-amino, mercaptoamino, thioalkylamino, acylamino, thioacylamino, cyano, carboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, thiocarboxylic acid or esters or thioesters or halides or anhydrides or amides thereof, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino and phenylhydrazino; optionally substituted heterocyclic radicals selected from the group consisting of oxabicycloheptyl, azabenzimidazolyl, azacycloheptyl, azacyclooctyl, azacyclononyl, azabicyclononyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydropyronyl, tetrahydroquinoleinyl,

tetrahydrothienyl and dioxide thereof, dihydrothienyl dioxide, dioxindolyl,
 dioxinyl, dioxenyl, dioxazinyl, thioxanyl, thioxolyl, thio-urazolyl, thiotriazolyl,
 thiopyranyl, thiopyronyl, coumarinyl, quinoleinyl, oxyquinoleinyl,
 quinuclidinyl, xanthinyl, dihydropyranyl, benzodihydrofuryl, benzothio-
 5 pyranyl, benzothiopyranyl, benzoxazinyl, benzoxazolyl, benzodioxolyl,
 benzodioxanyl, benzothiadiaazolyl, benzotriazinyl, benzothiazolyl,
 benzoxazolyl, phenothioxinyl, phenothiazolyl, phenothienyl, phenopyronyl,
 phenoxazolyl, pyridinyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl,
 thiomorpholinyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, tetrazinyl,
 10 triazolyl, benzotriazolyl, tetrazolyl, imidazolyl, pyrazolyl, thiazolyl,
 thiadiaazolyl, isothiazolyl, oxazolyl, oxadiaazolyl, pyrrolyl, furyl, dihydrofuryl,
 furoyl, hydantoinyl, dioxolanyl, dioxolyl, dithianyl, dithienyl, dithiinyl, thienyl,
 indolyl, indazolyl, benzofuryl, quinolyl, quinazolinyl, quinoxalinyl,
 carbazolyl, phenoxazinyl, phenothiazinyl, xanthenyl, purinyl, benzothienyl,
 15 naphthothienyl, thianthrenyl, pyranyl, pyronyl, benzopyronyl, isobenzofuranyl,
 chromenyl, phenoxathiinyl, indolizinyl, quinolizinyl, isoquinolyl,
 phthalazinyl, naphthiridinyl, cinnolyl, pteridinyl, carbolinyl, acridinyl,
 perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, imidazolyl,
 imidazolidinyl, benzimidazolyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl,
 20 pyrrolidinyl, piperazinyl, uridinyl, thymidinyl, cytidinyl, azirinyl, aziridinyl,
 diazirinyl, diaziridinyl, oxiranyl, oxaziridinyl, dioxiranyl, thiiranyl, azetyl,
 dihydroazetyl, azetidyl, oxetyl, oxetanyl, thietyl, thietanyl, diazabicyclo-
 octyl, diazetyl, diaziridinonyl, diaziridinethionyl, chromanyl, chromanonyl,
 thiochromanyl, thiochromanonyl, thiochromenyl, benzofuranyl,
 25 benzisothiazolyl, benzocarbazolyl, benzochromonyl, benziso-alloxazinyl,
 benzocoumarinyl, thiocoumarinyl, phenometoxazinyl, phenoparoxazinyl,
 phentriazinyl, thiodiazinyl, thiodiazolyl, indoxyl, thio-indoxyl, benzodiazinyl,
 phtalidyl, phtalimidinyl, phtalazonyl, alloxazinyl, xanthionyl, isatyl,
 isopyrazolyl, isopyrazolonyl, urazolyl, urazinyl, uretinyl, uretidinyl, succinyl,
 30 succinimido, benzylsultimyl and benzylsultamyl; aromatic or heterocyclic
 substituents substituted with an aliphatic spacer between the pteridine ring

- and the aromatic or heterocyclic substituent, whereby said aliphatic spacer is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms which may contain one or more functions, atoms or radicals selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, amino-acid, cyano, acylamino, thioacylamino, carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid or ester or thioester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; branched or straight, saturated or unsaturated aliphatic chains of 1 to 7 carbon atoms optionally containing one or more functions selected from the group consisting of carbonyl (oxo), thiocarbonyl, alcohol (hydroxyl), thiol, ether, thio-ether, acetal, thio-acetal, amino, imino, oximino, alkyloximino, aminoacid, cyano, acylamino; thioacylamino; carbamoyl, thiocarbamoyl, ureido, thio-ureido, carboxylic acid ester or halide or anhydride or amide, thiocarboxylic acid or ester or thioester or halide or anhydride or amide, nitro, thio C₁₋₇ alkyl, thio C₃₋₁₀ cycloalkyl, hydroxylamino, mercaptoamino, alkylamino, cycloalkylamino, alkenylamino, cycloalkenylamino, alkynylamino, arylamino, arylalkylamino, hydroxyalkylamino, mercaptoalkylamino, heterocyclic amino, hydrazino, alkylhydrazino, phenylhydrazino, sulfonyl, sulfinyl, sulfonamido and halogen; and
- R₃ is an atom or a group defined as R₄ or R₃ is selected from the group consisting of morpholinyl, amino, hydrogen, methyl, thiomethyl and chloro; or R₃ together with R₄ forms a homocyclic or heterocyclic radical;

and/or a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-*N*-oxide thereof and/or a solvate and/or a dihydro- or tetrahydropteridine derivative thereof.

5 21. A method of prevention or treatment according to claim 20, wherein an effective amount of the pharmaceutical composition corresponds to an amount in the range from 0.01 mg to 20 mg of the pteridine derivative per day and per kg body weight of the patient.

10 22. A method of prevention or treatment according to claim 20, wherein the said pharmaceutical composition further comprises one or more biologically-active drugs selected from the group consisting of immunosuppressant and/or immunomodulator drugs, antineoplastic drugs, and antiviral agents, or is administered in combination with an effective amount of a second pharmaceutical
15 composition comprising one or more biologically-active drugs selected from the group consisting of immunosuppressant and/or immunomodulator drugs, antineoplastic drugs, and antiviral agents.

20

25

30

ABSTRACT

This invention relates to a group of trisubstituted and tetrasubstituted pteridine derivatives, their pharmaceutically acceptable salts, N-oxides, solvates, dihydro- and tetrahydroderivatives and enantiomers, possessing unexpectedly desirable pharmaceutical properties, in particular which are highly active immunosuppressive agents, and as such are useful in the treatment in transplant rejection and/or in the treatment of certain inflammatory diseases. These compounds are also useful in preventing or treating cardiovascular disorders, allergic conditions, disorders of the central nervous system and cell proliferative disorders.

15

20

25

30


Figure 1


Figure 2


Figure 3

Figure 4





Figure 5



PATENT
ATTORNEY DOCKET NO. 50304/012001

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first, and sole inventor (if only one name is listed below) or an original, first, and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled **IMMUNOSUPPRESSIVE EFFECTS OF PTERIDINE DERIVATIVES**, the specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with 37 C.F.R. § 1.56.

FOREIGN PRIORITY RIGHTS: I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or (f), or § 365(b) of any foreign application(s) for patent, inventor's or plant breeder's rights certificate(s), or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below, and have also identified below any foreign application for patent, inventor's or plant breeder's rights certificate(s), or any PCT international application having a filing date before that of the application on which priority is claimed.

Country	Serial Number	Filing Date	Priority Claimed?
PCT	PCT/EP99/10320	December 28, 1999	Yes

PROVISIONAL PRIORITY RIGHTS: I hereby claim priority benefits under 35 U.S.C. § 119(e) of any United States provisional patent application(s) listed below filed by an inventor or inventors on the same subject matter as the present application and having a filing date before that of the application(s) of which priority is claimed:

Serial Number	Filing Date	Status
60/113,989	December 28, 1998	Abandoned

NON-PROVISIONAL PRIORITY RIGHTS: I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in 37 C.F.R. § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Serial Number	Filing Date	Status
09/869,468	October 10, 2001	Pending

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Paul T. Clark, Reg. No. 30,162; Karen L. Elbing, Ph.D., Reg. No. 35,238; Kristina Bieker-Brady, Ph.D., Reg. No. 39,109; Susan M. Michaud, Ph.D., Reg. No. 42,885; James D. DeCamp, Ph.D., Reg. No. 43,580; Jeffrey Ellison, Ph.D., Reg. No. 51,649; J. Cooper McDonald, Ph.D., Reg. No. 52,011; Jan Tittel, Ph.D., Reg. No. 52,290; Michael J. Belliveau, Ph.D., Reg. No. 52,608; R. Todd Armstrong, Ph.D., Reg. No. 54,590.

Address all telephone calls to: James D. DeCamp, at 617-428-0200.

Address all correspondence to: James D. DeCamp, at Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110. **Customer No. 21559**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001, and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Country of Citizenship
Mark Jozef Albert WAER	Heverlee BELGIUM	Waversebaan 263 B-3001 Heverlee BELGIUM	BELGIUM
Signature:			Date:

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Country of Citizenship
Piet André Maurits Maria HERDEWIJN	Rotselaar/Wezemaal BELGIUM	Olivierstraat 21 B-3111 Rotselaar/Wezemaal BELGIUM	BELGIUM
Signature:			Date:

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Country of Citizenship
Wolfgang Eugen PFLEIDERER	Konstanz GERMANY	Lindauerstrasse 47 D-78464 Konstanz GERMANY	GERMANY
Signature:			Date:

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/BE2004/000124

International filing date: 27 August 2004 (27.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 10/651,604
Filing date: 29 August 2003 (29.08.2003)

Date of receipt at the International Bureau: 17 March 2006 (17.03.2006)

Remark: Priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☒ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.